

REC'D 26 MAR 2003

V: PO

PCT

31 AUG 2004

Pl 977571

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

10/506455

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

March 18, 2003

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/361,948

FILING DATE: March 05, 2002

RELATED PCT APPLICATION NUMBER: PCT/US03/03120



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

P. Swain
P. SWAIN
Certifying Officer

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

03/05/02



1059 U.S.

03-01-02

Approved for filing through 04/11/98. OMB 0651-0037
Patent and Trademark Office, U.S. DEPARTMENT OF COMMERCE

Modified PTO/SB/16 (6-95)

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c)

Docket Number	P-15648	Type a plus sign (+) inside this box ->	+
---------------	---------	---	---

INVENTOR(s)/APPLICANT(s)			
LAST NAME	FIRST NAME	MIDDLE NAME	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)
Beals Kuchibhotla	John Uma	Michael	Indianapolis, Indiana Indianapolis, Indiana

TITLE OF THE INVENTION (280 characters max)
HETEROLOGOUS G-CSF FUSION PROTEINS

CORRESPONDENCE ADDRESS					
Eli Lilly and Company Patent Division/D.C. 1104 Lilly Corporate Center Indianapolis, Indiana 46285	 25885 PATENT TRADEMARK OFFICE				
STATE	IN	ZIP CODE	46285	COUNTRY	USA

ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification	Number of pages	53	<input type="checkbox"/> Small Entity Statement		
<input checked="" type="checkbox"/> Drawing(s)	Number of Sheets	1	<input type="checkbox"/> Other (Specify)		

METHOD OF PAYMENT (check one)		
<input type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees	PROVISIONAL FILING FEE AMOUNT (\$)	\$160.00
<input checked="" type="checkbox"/> The Assistant Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number:	05-0840	

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,
SIGNATURE Mark J. Stewart Date 03/05/02

TYPED or PRINTED NAME MARK J. STEWART REGISTRATION NO. (if appropriate) 43,936

☐ Additional inventors are being named on separately numbered sheets attached hereto

PROVISIONAL APPLICATION FOR PATENT FILING ONLY

"Express Mail" mailing label number <u>EL018702675US</u>	
Date of Deposit <u>March 5, 2002</u>	
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Arlington, VA, 22202.	
<u>Queen Thomas</u> Printed Name	<u>Queen Thomas</u> Signature

HETEROLOGOUS G-CSF FUSION PROTEINS**FIELD OF THE INVENTION**

5 The present invention relates to heterologous fusion
proteins, including analogs and derivatives thereof, fused
to proteins that have the effect of extending the *in vivo*
half-life of the proteins. These fusion proteins are
significant in human medicine, particularly in the treatment
of conditions treatable by stimulation of circulating
10 neutrophils, such as after chemotherapy regimens or in
chronic congenital neutropenia. More specifically, the
invention relates to novel heterologous fusion proteins with
granulocyte-colony stimulating factor activity.

BACKGROUND OF THE INVENTION

15 Among all blood cell lineages, the modulation of
neutrophil and platelet production has been of highest
interest to clinical oncologists and hematologists.
Myelosuppression is the single most severe complication of
20 cancer chemotherapy, and a major cause of treatment delay
during multiple-cycle or combination chemotherapy. It is
also the major dose-limiting factor for most
chemotherapeutic agents. Due to the short half-lives of
neutrophils in peripheral blood, life-threatening falls in
25

"Express Mail" mailing label number EL018702675US

Date of Deposit March 5, 2002

30 I hereby certify that this paper or fee is being deposited with the United
States Postal Service "Express Mail Post Office to Addressee" service under 37
C.F.R. 1.10 on the date indicated above and is addressed to the Assistant
Commissioner for Patents, Washington, D.C. 20231.

35 Queen Adams
Printed Name

Queen Adams
Signature

neutrophil levels are seen after a number of conventional anti-tumor chemotherapy regimens.

The most prominent regulator of granulopoiesis is granulocyte-colony stimulating factor (G-CSF). G-CSF induces proliferation and differentiation of hematopoietic progenitor cells resulting in increased numbers of circulating neutrophils. G-CSF also stimulates the release of mature neutrophils from bone marrow and activates their functional state. [Souza L.M., et al. (1986) *Science* 232:61-65]. Thus, therapeutic proteins with G-CSF activity have tremendous value in situations where there are reduced circulating levels of neutrophilic granulocytes.

However, the usefulness of therapy using G-CSF peptides has been limited by their short plasma half-life. Thus, they must be administered intravenously or subcutaneously at fairly frequent intervals (once or twice a day) in order to maintain their neutrophil stimulating properties. In addition, this short half-life limits the performance of the drug to traditional drug delivery systems. It would clearly benefit the treatment of patients with abnormally low neutrophils, and reduce the discomfort and inconvenience associated with frequent injections to provide a pharmaceutical agent that could be administered less frequently and optionally by alternative routes of administration. Thus, a need exists to develop agents that stimulate the production of mature neutrophils and are more optimal in their duration of effect.

The present invention overcomes the problems associated with delivering a compound that has a short plasma half-life in two respects. First, G-CSF is hyperglycosylated. The carbohydrate content of G-CSF is altered by substituting amino acids that can act as substrates for glycosylating enzymes in mammalian cells. Most significantly, the present invention encompasses fusion of these hyperglycosylated G-

BRIEF SUMMARY OF THE INVENTION

10

15

- 20

- a glycine rich peptide;
- a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]_n where n is 1, 2, 3, 4, or 5; and
- a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]₃

25

30

	1				5					10					15	
	Thr	Pro	Leu	Gly	Pro	Ala	Ser	Ser	Leu	Pro	Gln	Ser	Phe	Leu	Leu	Lys
				20					25					30		
35	Xaa	Leu	Glu	Gln	Val	Arg	Lys	Ile	Gln	Gly	Asp	Gly	Ala	Ala	Leu	Gln
			35					40					45			
	Glu	Lys	Leu	Cys	Xaa	Xaa	Xaa	Lys	Leu	Cys	His	Pro	Glu	Glu	Leu	Val
		50					55					60				
	Leu	Leu	Gly	His	Ser	Leu	Gly	Ile	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa

	65				70				75				80
	Xaa	Xaa	Xaa	Xaa	Xaa	Gln	Leu	Ala	Gly	Cys	Leu	Ser	Gln
					85				90				95
5	Gly	Leu	Phe	Leu	Tyr	Gln	Gly	Leu	Leu	Gln	Ala	Leu	Xaa
				100					105				110
	Xaa	Glu	Leu	Gly	Pro	Thr	Leu	Asp	Thr	Leu	Gln	Leu	Asp
				115					120				125
	Phe	Ala	Thr	Thr	Ile	Trp	Gln	Gln	Met	Glu	Glu	Leu	Gly
				130					135				140
10	Ala	Leu	Gln	Pro	Xaa	Xaa	Xaa	Ala	Met	Pro	Ala	Phe	Xaa
						150							160
	Gln	Arg	Arg	Ala	Gly	Gly	Val	Leu	Val	Ala	Ser	His	Leu
					165					170			
	Leu	Glu	Val	Ser	Tyr	Arg	Val	Leu	Arg	His	Leu	Ala	Gln
15													Pro

(I)

wherein:

- Xaa at position 17 is Cys, Ala, Leu, Ser, or Glu;
 Xaa at position 37 is Ala or Asn;
 Xaa at position 38 is Thr, or any other amino acid except
 20 Pro;
 Xaa at position 39 is Tyr, Thr, or Ser;
 Xaa at position 57 is Pro or Val;
 Xaa at position 58 is Trp or Asn;
 Xaa at position 59 is Ala or any other amino acid except
 25 Pro;
 Xaa at position 60 is Pro, Thr, Asn, or Ser,
 Xaa at position 61 is Leu, or any other amino acid except
 Pro;
 Xaa at position 62 is Ser or Thr;
 30 Xaa at position 63 is Ser or Asn;
 Xaa at position 64 is Cys or any other amino acid except
 Pro;
 Xaa at position 65 is Pro, Ser, or Thr;
 Xaa at position 66 is Ser or Thr;
 35 Xaa at position 67 is Gln or Asn;
 Xaa at position 68 is Ala or any other amino acid except
 Pro;
 Xaa at position 69 is Leu, Thr, or Ser
 Xaa at position 93 is Glu or Asn
 40 Xaa at position 94 is Gly or any other amino acid except
 Pro;
 Xaa at position 95 is Ile, Asn, Ser, or Thr;
 Xaa at position 97 is Pro, Ser, Thr, or Asn;
 Xaa at position 133 is Thr or Asn;
 45 Xaa at position 134 is Gln or any other amino acid except
 Pro;
 Xaa at position 135 is Gly, Ser, or Thr
 Xaa at position 141 is Ala or Asn;
 Xaa at position 142 is Ser or any other amino acid except
 50 Pro; and

Xaa at position 143 is Ala, Ser, or Thr;

and wherein:

5 Xaa at positions 37, 38, and 39 constitute region 1;
Xaa at positions 58, 59, and 60 constitute region 2;
Xaa at positions 59, 60, and 61 constitute region 3;
Xaa at positions 60, 61, and 62 constitute region 4;
Xaa at positions 61, 62, and 63 constitute region 5;
Xaa at positions 62, 63, and 64 constitute region 6;
10 Xaa at positions 63, 64, and 65 constitute region 7;
Xaa at positions 64, 65, and 66 constitute region 8;
Xaa at positions 67, 68, and 69 constitute region 9;
Xaa at positions 93, 94, and 95 constitute region 10;
Xaa at positions 94, 95, and Ser at position 96
15 constitute region 11;
Xaa at positions 95, and 97, and Ser at position 96
constitute region 12;
Xaa at positions 133, 134, and 135 constitute
region 13;
20 Xaa at positions 141, 142, and 143 constitute
region 14;

and provided that at least one of regions 1 through 14
comprises the sequence Asn Xaa1 Xaa2 wherein Xaa1 is any
25 amino acid except Pro and Xaa2 is Ser or Thr.

Thus, the heterologous fusion proteins of the present
invention include analogs wherein one or any combination of
two or more regions comprise the sequence Asn Xaa1 Xaa2
wherein Xaa1 is any amino acid except Pro and Xaa2 is Ser or
30 Thr.

Preferred hyperglycosylated G-CSF analogs that make up
part of the heterologous fusion proteins of the present
invention, include the following:

35 a) G-CSF[A37N,Y39T]

- b) G-CSF[P57V,W58N,P60T]
c) G-CSF[P60N,S62T]
d) G-CSF[S63N,P65T]
e) G-CSF[Q67N,L69T]
5 f) G-CSF[E93N,I95T]
g) G-CSF[T133N,G135T]
h) G-CSF[A141N,A143T]
i) G-CSF[A37N,Y39T,P57V,W58N,P60T]
j) G-CSF[A37N,Y39T,P60N,S62T]
10 k) G-CSF[A37N,Y39T,S63N,P65T]
l) G-CSF[A37N,Y39T,Q67N,L69T]
m) G-CSF[A37N,Y39T,E93N,I95T]
n) G-CSF[A37N,Y39T,T133N,G135T]
o) G-CSF[A37N,Y39T,A141N,A143T]
15 p) G-CSF[A37N,Y39T,P57V,W58N,P60T,S63N,P65T]
q) G-CSF[A37N,Y39T,P57V,W58N,P60T,Q67N,L69T]
r) G-CSF[A37N,Y39T,S63N,P65T,E93N,I95T]

The present invention also includes heterologous fusion proteins, which are the product of the expression in a host cell of an exogenous DNA sequence, which comprises a DNA sequence encoding a heterologous fusion protein of Formula I (described above) fused to a DNA sequence encoding human albumin or the Fc portion of an immunoglobulin.

A hyperglycosylated heterologous fusion protein of the present invention also includes polynucleotides encoding the heterologous fusion protein described herein, vectors comprising these polynucleotides and host cells transfected or transformed with the vectors described herein. Also included is a process for producing a heterologous fusion protein comprising the steps of transcribing and translating a polynucleotide described herein under conditions wherein the heterologous fusion protein is expressed in detectable amounts.

The present invention encompasses a method for increasing neutrophil levels in a mammal comprising the administration of a therapeutically effective amount of a heterologous fusion protein described above. The present invention also includes the use of the heterologous fusion proteins described above for the manufacture of a medicament for the treatment of patients with insufficient circulating neutrophil levels.

The present invention also encompasses a pharmaceutical formulation adapted for the treatment of patients with insufficient neutrophil levels comprising a glycosylated protein as described above.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention is further illustrated with reference to the following drawings:

Figure 1: Schematic illustrating fourteen regions in human G-CSF wherein the amino acid sequence can be mutated to create functional glycosylation sites.

DETAILED DESCRIPTION OF THE INVENTION

The present invention comprises heterologous fusion proteins. As used herein, the term heterologous fusion protein means a hyperglycosylated G-CSF analog fused to human albumin, a human albumin analog, a human albumin fragment, the Fc portion of an immunoglobulin, an analog of the Fc portion of an immunoglobulin, or a fragment of the Fc portion of an immunoglobulin. The G-CSF analog may be fused directly, or fused via a peptide linker, to an albumin or Fc protein. The albumin and Fc portion may be fused to the G-CSF analogs at either terminus or at both termini. These heterologous fusion proteins are biologically active and have an increased half-life compared to native G-CSF.

Hyperglycosylated G-CSF Analogs

Hyperglycosylated analogs of G-CSF as used herein refers to analogs of G-CSF with one or more changes in the amino acid sequence which result in an increase in the number of sites for carbohydrate attachment compared with native human G-CSF expressed in animal cells *in vivo*. In addition, hyperglycosylated G-CSF analogs include human G-CSF wherein the O-linked glycosylation site at position 133 is replaced with an N-linked glycosylation site. Analogs are generated by site directed mutagenesis having substitution of amino acid residues creating new sites that are available for glycosylation. G-CSF analogs having a greater carbohydrate content than that found in native human G-CSF are generated by adding glycosylation sites that do not perturb the secondary, tertiary, and quaternary structure required for activity. The heterologous fusion proteins of the present invention thus have a larger mass and an increased negative charge compared to native G-CSF, they will not be as rapidly cleared from the circulation.

It is preferred that the hyperglycosylated G-CSF analog portion of the heterologous fusion protein have 1, 2, 3, or 4 additional sites for N-glycosylation. Figure 1 illustrates fourteen different regions that can be glycosylated with very little effect on *in vitro* activity. Each region may be mutated to the consensus site for N-glycosylation addition which is Asn X1 X2 wherein X1 is any amino acid except Pro and X2 is Ser or Thr. It is preferred that the X1 amino acid be any other amino acid except Trp, Asp, Glu, or Leu and it is most preferred that the X1 amino acid be the naturally occurring amino acid. The scope of the present invention includes analogs wherein a single region (1 through 14) is mutated or wherein a region is mutated in combination with one or more other regions.

A representative number of fusion proteins having single of multiple additional glycosylation sites have been expressed, purified, and characterized. The mutated amino

acids in the hyperglycosylated G-CSF portion of the heterologous fusion protein are identified in parentheses. For example, G-CSF[A37N,Y39T] is G-CSF wherein the amino acids at positions 37 and 39 have been substituted to create a glycosylation site. This site of carbohydrate attachment is illustrated as region 1 in Figure 1. G-CSF[A37N,Y39T,P57V,W58N,P60T] is an example of a G-CSF analog wherein amino acids in region 1 and region 2 are mutated to provide two functional glycosylation sites on a single molecule (Figure 1).

G-CSF[A37N,Y39T,P57V,W58N,P60T,Q67N,L69T] is an example of a G-CSF analog wherein the amino acids in region 1, region 2, and region 9 are mutated to provide three functional glycosylation sites on a single molecule (Figure 1).

Native G-CSF can be used as the backbone to create the hyperglycosylated G-CSF analog portion of a particular heterologous fusion protein. In addition, the native G-CSF backbone can be modified such that substitutions in the regions defined in Figure 1 are made in the context of a different or improved heterologous fusion protein. For example, native G-CSF with a Cysteine to Alanine substitution at position 17 may reduce aggregation and enhance stability and thus, can be used as the backbone used to create the hyperglycosylated G-CSF analog portion of the heterologous fusion proteins of the present invention.

In addition, Reidhaar-Olson, et al., through alanine scanning mutagenesis, describe residues critical to the activity of human G-CSF. [Reidhaar-Olson, et al. (1996) *Biochemistry* 35:9034-9041; See also Young, et al. (1997) *Protein Science* 6:1228-1236]. Thus, the hyperglycosylated G-CSF analogs can be modified by substituting amino acids outside the glycosylated regions described in Figure 1.

The heterologous fusion proteins of the present invention also encompass G-CSF analogs wherein the O-linked glycosylation site at position 133 is mutated to serve as an N-linked glycosylation site. The N-linked carbohydrate will generally have a higher sialic acid content which will protect it from the rapid clearance mechanisms associated with native G-CSF.

The functions of a carbohydrate chain greatly depend on the structure of the attached carbohydrate moiety.

Typically compounds with a higher sialic acid content will have better stability and longer half-lives *in vivo*. The N-linked oligosaccharides contain sialic acid in both an $\alpha 2,3$ and an $\alpha 2,6$ linkage to galactose. [Takeuchi, et al. (1988) *J. Biol. Chem.* 263:3657]. Typically the sialic acid in the $\alpha 2,3$ linkage is added to galactose on the mannose $\alpha 1,6$ branch and the sialic acid in the $\alpha 2,6$ linkage is added to the galactose on the mannose $\alpha 1,3$ branch. The enzymes that add these sialic acids (β -galactoside $\alpha 2,3$ sialyltransferase and β -galactoside $\alpha 2,6$ sialyltransferase) are most efficient at adding sialic acid to the mannose $\alpha 1,6$ and mannose $\alpha 1,3$ branches respectively.

Tetra-antennary N-linked oligosaccharides most commonly provide four possible sites for sialic acid attachment while bi- and tri-antennary oligosaccharide chains, which can substitute for the tetra-antennary form at Asn-linked sites, commonly have at most only two or three sialic acids attached. O-linked oligosaccharides commonly provide only two sites for sialic acid attachment. Mammalian cell cultures can be screened for those cells that preferentially add tetra-antennary chains to the G-CSF analogs of the present invention, thereby maximizing the number of sites for sialic acid attachment. Different types of mammalian cells also differ with respect to the transferase enzymes present and consequently the sialic acid content and type of oligosaccharide attached at each site. One way to optimize the carbohydrate content for a given G-CSF analog is to

express the analog in a cell line wherein an expression plasmid containing DNA encoding a specific sialyl transferase (e.g., $\alpha 2,6$ sialyltransferase) is co-transfected with the G-CSF analog expression plasmid.

- 5 Alternatively a host cell line may be stably transfected with a sialyl transferase cDNA and that host cell used to express the G-CSF analog of interest. Thus, it is preferable if the oligosaccharide structure and sialic acid content are optimized for each analog encompassed by the
10 present invention.

Heterologous Fc fusion proteins:

- 15 The hyperglycosylated G-CSF analogs described above can be fused directly or via a peptide linker to the Fc portion of an immunoglobulin. For example SEQ ID NO:23 represents the sequence of a G-CSF-IgG1 Fc fusion protein, SEQ ID NO:24 represents the sequence of a G-CSF-IgG4 Fc fusion protein, and SEQ ID NO:25 represents the sequence of a G-CSF-human albumin fusion protein.

- 20 Immunoglobulins are molecules containing polypeptide chains held together by disulfide bonds typically having two light chains and two heavy chains. In each chain, one domain (V) has a variable amino acid sequence depending on the antibody specificity of the molecule. The other domains
25 (C) have a rather constant sequence common to molecules of the same class.

- As used herein, the Fc portion of an immunoglobulin has the meaning commonly given to the term in the field of immunology. Specifically, this term refers to an antibody
30 fragment which is obtained by removing the two antigen binding regions (the Fab fragments) from the antibody. Thus, the Fc portion is formed from approximately equal sized fragments of the constant region from both heavy chains, which associate through non-covalent interactions
35 and disulfide bonds. The Fc portion can include the hinge regions and extend through the CH2 and CH3 domains to the C-

terminus of the antibody. For example, SEQ ID NOs: 18 and 19 represent the sequence of a Fc region encompassing the hinge region as well as the CH2 and CH3 domains. Other representative hinge regions for human and mouse

5 immunoglobulins can be found in Antibody Engineering, A Practical Guide, Borrebaeck, C.A.K., ed., W.H. Freeman and Co., 1992, the teachings of which are herein incorporated by reference.

10 There are five types of human immunoglobulin Fc regions with different effector and pharmacokinetic properties: IgG, IgA, IgM, IgD, and IgE. IgG is the most abundant immunoglobulin in serum. IgG also has the longest half-life in serum of any immunoglobulin (23 days). Unlike other immunoglobulins, IgG is efficiently recirculated following
15 binding to an Fc receptor. There are four IgG subclasses G1, G2, G3, and G4, each of which have different effector functions. G1, G2, and G3 can bind C1q and fix complement while G4 cannot. Even though G3 is able to bind C1q more efficiently than G1, G1 is more effective at mediating
20 complement-directed cell lysis. G2 fixes complement very inefficiently. The C1q binding site in IgG is located at the carboxy terminal region of the CH2 domain.

All IgG subclasses are capable of binding to Fc receptors (CD16, CD32, CD64) with G1 and G3 being more
25 effective than G2 and G4. The Fc receptor-binding region of IgG is formed by residues located in both the hinge and the carboxy terminal regions of the CH2 domain.

IgA can exist both in a monomeric and dimeric form held together by a J-chain. IgA is the second most abundant Ig
30 in serum, but it has a half-life of only 6 days. IgA has three effector functions. It binds to an IgA specific receptor on macrophages and eosinophils, which drives phagocytosis and degranulation, respectively. It can also fix complement via an unknown alternative pathway.

35 IgM is expressed as either a pentamer or a hexamer, both of which are held together by a J-chain. IgM has a

serum half-life of 5 days. It binds weakly to C1q via a binding site located in its CH3 domain. IgD has a half-life of 3 days in serum. It is unclear what effector functions are attributable to this Ig. IgE is a monomeric Ig and has
5 a serum half-life of 2.5 days. IgE binds to two Fc receptors which drives degranulation and results in the release of pro-inflammatory agents.

Depending on the desired *in vivo* effect, the heterologous fusion proteins of the present invention may
10 contain any of the isotypes described above or may contain mutated Fc regions wherein the complement and/or Fc receptor binding functions have been altered. For example, one embodiment of the present invention is a heterologous fusion protein wherein the Fc portion comprises a human IgG4
15 sequence wherein serine at position 229 is changed to proline which reduces the effector function of the protein. (See SEQ ID NO: 22).

The heterologous fusion proteins of the present invention may contain the entire Fc portion of an
20 immunoglobulin, fragments of the Fc portion of an immunoglobulin, or analogs thereof fused to a G-CSF analog. Furthermore, the Fc portion may be fused at either terminus or at both termini.

The heterologous fusion proteins of the present invention can consist of single chain proteins or as multi-chain polypeptides. Two or more Fc fusion proteins can be
25 produced such that they interact through disulfide bonds that naturally form between Fc regions. These multimers can be homogeneous with respect to the G-CSF analog or they may
30 contain different G-CSF analogs fused at the N-terminus of the Fc portion of the fusion protein.

Regardless of the final structure of the fusion protein, the Fc or Fc-like region serves to prolong the *in vivo* plasma half-life of the G-CSF analog compared to native
35 G-CSF. Furthermore, the fused G-CSF analog must retain some

biological activity. Biological activity can be determined by in vitro and in vivo methods known in the art.

Since the Fc region of IgG produced by proteolysis has the same in vivo half-life as the intact IgG molecule and

5 Fab fragments are rapidly degraded, it is believed that the relevant sequence for prolonging half-life resides in the CH2 and/or CH3 domains. Further, it has been shown in the literature that the catabolic rates of IgG variants that do not bind the high-affinity Fc receptor or C1q are

10 indistinguishable from the rate of clearance of the parent wild-type antibody, indicating that the catabolic site is distinct from the sites involved in Fc receptor or C1q binding. [Wawrzynczak et al., (1992) *Molecular Immunology* 29:221]. Site-directed mutagenesis studies using a murine
15 IgG1 Fc region suggested that the site of the IgG1 Fc region that controls the catabolic rate is located at the CH2-CH3 domain interface.

Based on these studies, Fc regions can be modified at the catabolic site to optimize the half-life of the fusion
20 proteins. It is preferable that the Fc region used for the heterologous fusion proteins of the present invention be derived from an IgG1 such as that represented by SEQ ID NO: 21 or an IgG4 Fc region such as that represented by SEQ ID NO: 22. It is even more preferable that the Fc region be
25 IgG4 or derived from IgG4. Preferably the IgG Fc region contains both the CH2 and CH3 regions including the hinge region.

Heterologous albumin fusion proteins:

30 The G-CSF analogs described above can be fused directly or via a peptide linker to albumin or an analog, fragment, or derivative thereof. SEQ ID NO:25 provides a representative G-CSF-human albumin fusion protein.

Generally the albumin proteins making up part of the
35 heterologous fusion proteins of the present invention can be derived from albumin cloned from any species. However,

human albumin and fragments and analogs thereof are preferred to reduce the risk of the fusion protein being immunogenic in humans. Human serum albumin (HA) consists of a single non-glycosylated polypeptide chain of 585 amino acids with a formula molecular weight of 66,500. The amino acid sequence of human albumin is represented as SEQ ID NO: 20. [See Meloun, et al. (1975) FEBS Letters 58:136; Behrens, et al. (1975) Fed. Proc. 34:591; Lawn, et al. (1981) Nucleic Acids Research 9:6102-6114; Minghetti, et al. (1986) J. Biol. Chem. 261:6747]. A variety of polymorphic variants as well as analogs and fragments of albumin have been described. [See Weitkamp, et al., (1973) Ann. Hum. Genet. 37:219]. For example, in EP 322,094, the inventors disclose various shorter forms of HA. Some of these fragments include HA(1-373), HA(1-388), HA(1-389), HA(1-369), and HA(1-419) and fragments between 1-369 and 1-419. EP 399,666 discloses albumin fragments that include HA(1-177) and HA(1-200) and fragments between HA(1-177) and HA(1-200).

It is understood that the heterologous fusion proteins of the present invention include hyperglycosylated G-CSF analogs that are coupled to any albumin protein including fragments, analogs, and derivatives wherein such fusion protein is biologically active and has a longer plasma half-life than native G-CSF. Thus, the albumin portion of the fusion protein need not necessarily have a plasma half-life equal to that of native human albumin. In addition, the albumin may be fused to either terminus or both termini of the hyperglycosylated G-CSF analog.

The heterologous fusion proteins of the present invention encompass proteins having conservative amino acid substitutions in the hyperglycosylated G-CSF analog and/or the Fc or albumin portion of the fusion protein.

The term "amino acid" is used herein in its broadest sense, and includes naturally occurring amino acids as well as non-naturally occurring amino acids, including amino acid

analogs and derivatives. The latter includes molecules containing an amino acid moiety. One skilled in the art will recognize, in view of this broad definition, that reference herein to an amino acid includes, for example, naturally occurring proteogenic L-amino acids; D-amino acids; chemically modified amino acids such as amino acid analogs and derivatives; naturally occurring non-proteogenic amino acids such as norleucine, β -alanine, ornithine, GABA, etc.; and chemically synthesized compounds having properties known in the art to be characteristic of amino acids. As used herein, the term "proteogenic" indicates that the amino acid can be incorporated into a peptide, polypeptide, or protein in a cell through a metabolic pathway.

The incorporation of non-natural amino acids, including synthetic non-native amino acids, substituted amino acids, or one or more D-amino acids into the heterologous fusion proteins of the present invention can be advantageous in a number of different ways. D-amino acid-containing peptides, etc., exhibit increased stability *in vitro* or *in vivo* compared to L-amino acid-containing counterparts. Thus, the construction of peptides, etc., incorporating D-amino acids can be particularly useful when greater intracellular stability is desired or required. More specifically, D-peptides, etc., are resistant to endogenous peptidases and proteases, thereby providing improved bioavailability of the molecule, and prolonged lifetimes *in vivo* when such properties are desirable. Additionally, D-peptides, etc., cannot be processed efficiently for major histocompatibility complex class II-restricted presentation to T helper cells, and are therefore less likely to induce humoral immune responses in the whole organism.

Although the heterologous fusion proteins of the present invention can be made by a variety of different methods, recombinant methods are preferred.

Construction of DNA encoding the heterologous fusion proteins of the present invention:

Wild type albumin and immunoglobulin proteins can be obtained from a variety of sources. For example, these proteins can be obtained from a cDNA library prepared from tissue or cells which express the mRNA of interest at a detectable level. Libraries can be screened with probes designed using the published DNA or protein sequence for the particular protein of interest.

Screening a cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, NY (1989). An alternative means to isolate a gene encoding an albumin or immunoglobulin protein is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., *PCR Primer: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, NY (1995)]. PCR primers can be designed based on published sequences.

Generally the full-length wild-type sequences cloned from a particular species can serve as a template to create analogs, fragments, and derivatives that retain the ability to confer a longer plasma half-life on the G-CSF analog that is part of the fusion protein. It is preferred that the Fc and albumin portions of the heterologous fusion proteins of the present invention be derived from the native human sequence in order to reduce the risk of potential immunogenicity of the fusion protein in humans.

In particular, it is preferred that the immunoglobulin portion of a fusion protein encompassed by the present invention contain only an Fc fragment of the immunoglobulin. Depending on whether particular effector functions are desired and the structural characteristics of the fusion protein, an Fc fragment may contain the hinge region along with the CH2 and CH3 domains or some other combination thereof. These Fc fragments can be generated using PCR

techniques with primers designed to hybridize to sequences corresponding to the desired ends of the fragment. Similarly, if fragments of albumin are desired, PCR primers can be designed which are complementary to internal albumin sequences. PCR primers can also be designed to create restriction enzyme sites to facilitate cloning into expression vectors.

DNA encoding human G-CSF can be obtained from a cDNA library prepared from tissue or cells which express G-CSF mRNA at a detectable level such as monocytes, macrophages, vascular endothelial cells, fibroblasts, and some human malignant and leukemic myeloblastic cells. Libraries can be screened with probes designed using the published DNA sequence for human G-CSF. [Souza L., et al. (1986) *Science* 232:61-65]. Screening a cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, NY (1989). An alternative means to isolate the gene encoding human G-CSF is to use PCR methodology [Sambrook, et al., supra; Dieffenbach, et al., *PCR Primer: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, NY (1995)].

The glycosylated G-CSF analogs of the present invention can be constructed by a variety of mutagenesis techniques well known in the art. Specifically, a representative number of glycosylated G-CSF analogs were constructed using mutagenic PCR from a cloned wild-type human G-CSF DNA template (Example 1).

The glycosylated G-CSF analogs of the present invention may be produced by other methods including recombinant DNA technology or well known chemical procedures, such as solution or solid-phase peptide synthesis, or semi-synthesis in solution beginning with protein fragments coupled through conventional solution methods.

Recombinant DNA methods are preferred for producing the glycosylated G-CSF analogs of the present invention. Host

cells are transfected or transformed with expression or cloning vectors described herein for glycosylated G-CSF analog production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences (Example 2). The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation

Physical stability is an essential feature for therapeutic formulations. The physical stability of the heterologous fusion proteins of the present invention depends on their conformational stability, the number of charged residues (pI of the protein), the ionic strength and pH of the formulation, and the protein concentration, among other possible factors. As discussed previously, the G-CSF analog portion of the heterologous fusion proteins can be successfully glycosylated and expressed such that the three dimensional structure is maintained. Because these heterologous fusion proteins are able to fold properly in a hyperglycosylated state, they will have improved conformational and physical stability relative to wild-type G-CSF.

While wild-type G-CSF produced in mammalian cells and bacterial cells has similar activity *in vivo*, the mammalian cell-produced protein has increased conformational and physical stability due to the presence of a single O-linked sugar moiety present at position 133. Thus, the G-CSF analog portion of the heterologous fusion proteins, which has an increased glycosylation content compared to wild-type G-CSF produced in mammalian or bacterial cells, will confer increased stability on the heterologous fusion protein. Furthermore, it is likely that glycosylation may inhibit inter-domain interactions and consequently enhance stability by preventing inter-domain disulfide shuffling.

The gene encoding a heterologous fusion protein can be constructed by ligating DNA encoding a G-CSF analog in-frame

to DNA encoding an albumin or Fc protein. The gene encoding the G-CSF analog and the gene encoding the albumin or Fc protein can also be joined in-frame via DNA encoding a linker peptide.

5 The *in vivo* function and stability of the heterologous fusion proteins of the present invention can be optimized by adding small peptide linkers to prevent potentially unwanted domain interactions. Although these linkers can potentially be any length and consist of any combination of amino acids,
10 it is preferred that the length be no longer than necessary to prevent unwanted domain interactions and/or optimize biological activity and/or stability. Generally, the linkers should not contain amino acids with extremely bulky side chains or amino acids likely to introduce significant
15 secondary structure. It is preferred that the linker be serine-glycine rich and be less than 30 amino acids in length. It is more preferred that the linker be no more than 20 amino acids in length. It is even more preferred that the linker be no more than 15 amino acids in length. A
20 preferred linker contains repeats of the sequence Gly-Gly-Gly-Gly-Ser. It is preferred that there be between 2 and 6 repeats of this sequence. It is even more preferred that there be between 3 and 4 repeats of this sequence.

 To construct the heterologous G-CSF fusion proteins, the
25 DNA encoding wild-type G-CSF, albumin, and Fc polypeptides and fragments thereof can be mutated either before ligation or in the context of a cDNA encoding an entire fusion protein. A variety of mutagenesis techniques are well known in the art. For example, a mutagenic PCR method utilizes
30 strand overlap extension to create specific base mutations for the purposes of changing a specific amino acid sequence in the corresponding protein. This PCR mutagenesis requires the use of four primers, two in the forward orientation (primers A and C) and two in the reverse orientation
35 (primers B and D). A mutated gene is amplified from the wild-type template in two different stages. The first

reaction amplifies the gene in halves by performing an A to B reaction and a separate C to D reaction wherein the B and C primers target the area of the gene to be mutated. When aligning these primers with the target area, they contain mismatches for the bases that are targeted to be changed. Once the A to B and C to D reactions are complete, the reaction products are isolated and mixed for use as the template for the A to D reaction. This reaction then yields the full, mutated product.

Once a gene encoding an entire fusion protein is produced it can be cloned into an appropriate expression vector. Specific strategies that can be employed to make the G-CSF fusion proteins of the present invention are described in example 1.

General methods to recombinantly express the heterologous fusion proteins of the present invention:

Host cells are transfected or transformed with expression or cloning vectors described herein for heterologous fusion protein production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in *Mammalian Cell Biotechnology: A Practical Approach*, M. Butler, ed. (IRL Press, 1991) and Sambrook, et al., supra. Methods of transfection are known to the ordinarily skilled artisan, for example, CaPO_4 and electroporation. General aspects of mammalian cell host system transformations have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of van Solingen, et al., *J Bact.* 130(2): 946-7 (1977) and Hsiao, et al., *Proc. Natl. Acad. Sci. USA* 76(8): 3829-33 (1979). Suitable host

cells for the expression of the fusion proteins of the present invention are derived from multicellular organisms.

The fusion proteins of the present invention may be recombinantly produced directly, or as a protein having a signal sequence or other additional sequences which create a specific cleavage site at the N-terminus of the mature fusion protein. In general, the signal sequence may be a component of the vector, or it may be a part of the fusion protein-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* cc-factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179), or the signal described in WO 90/13646. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species as well as viral secretory leaders.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the fusion protein-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described [Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77(7): 4216-20 (1980)]. A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid Yrp7 [Stinchcomb, et al., *Nature* 282(5734): 39-43 (1979); Kingsman, et al., *Gene* 7(2): 141-52 (1979); Tschumper, et al., *Gene* 10(2): 157-66 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in

tryptophan, for example, ATCC No. 44076 or PEPC1 [Jones, *Genetics* 85: 23-33 (1977)].

Various forms of a fusion protein may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g., Triton-X 100) or by enzymatic cleavage. Cells employed in expression of a fusion protein can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

Purification of the heterologous fusion proteins of the present invention:

Once the heterologous fusion proteins of the present invention are expressed in the appropriate host cell, the analogs can be isolated and purified. The following procedures are exemplary of suitable purification procedures:

Various methods of protein purification may be employed and such methods are known in the art and described, for example, in Deutscher, *Methods in Enzymology* 182: 83-9 (1990) and Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag, NY (1982). The purification step(s) selected will depend on the nature of the production process used and the particular fusion protein produced. For example, fusion proteins comprising an Fc fragment can be effectively purified using a Protein A or Protein G affinity matrix. Low or high pH buffers can be used to elute the fusion protein from the affinity matrix. Mild elution conditions will aid in preventing irreversible denaturation of the fusion protein. Imidazole-containing buffers can also be used. Example 3 describes some successful purification protocols for the fusion proteins of the present invention.

Characterization of the heterologous fusion proteins of the present invention:

Numerous methods exist to characterize the fusion proteins of the present invention. Some of these methods include: SDS-PAGE coupled with protein staining methods or immunoblotting using anti-IgG, anti-HA and anti-G-CSF antibodies. Other methods include matrix assisted laser desorption/ionization-mass spectrometry (MALDI-MS), liquid chromatography/mass spectrometry, isoelectric focusing, analytical anion exchange, chromatofocussing, and circular dichroism to name a few. A representative number of heterologous fusion proteins were characterized using SDS-PAGE coupled with immunoblotting as well as mass spectrometry

For example, Table 2 illustrates the calculated molecular mass for a representative number of fusion proteins as well as the observed mass (as measured by protease mapping/LC-MS). The relative differences between observed mass and mass calculated for a nonglycosylated protein are indicative of the extent of glycosylation.

Table 2

Construct	Theoretical mass (kDa)	Observed mass (kDa)
C17A (GCSF)-IgG1 Fc	89.4	93.0
C17A (GCSF)-IgG4 Fc	88.7	92.7
C17A (GCSF)-HSA	86.0	85.1
C17A, A37N, S63N, E93N (GCSF)-IgG1 Fc	89.4	97-114 (104.7)
C17A, A37N, S63N, E93N (GCSF)-IgG4 Fc	88.7	96-112 (103.8)
C17A, A37N, S63N, E93N (GCSF)-HSA	86.0	87-98 (92.5)
C17A, A37N, W58N, Q67N (GCSF)-IgG1 Fc	89.3	95-110 (103.8)
C17A, A37N, W58N, Q67N (GCSF)-IgG4 Fc	88.8	95-110 (102.4)
C17A, A37N, W58N, Q67N (GCSF)-HSA	85.9	87-98 (93.1)

Administration and Preparation of Compositions:

The heterologous fusion proteins of the present invention may be formulated with one or more excipients. The active

fusion proteins of the present invention may be combined with a pharmaceutically acceptable buffer, and the pH adjusted to provide acceptable stability, and a pH acceptable for administration such as parenteral administration.

5 Optionally, one or more pharmaceutically-acceptable anti-microbial agents may be added. Meta-cresol and phenol are preferred pharmaceutically-acceptable microbial agents. One or more pharmaceutically-acceptable salts may be added to adjust the ionic strength or tonicity. One or more excipients
10 may be added to adjust the isotonicity of the formulation. Glycerin is an example of an isotonicity-adjusting excipient. Pharmaceutically acceptable means suitable for administration to a human or other animal and thus, does not contain toxic elements or undersirable contaminants and does not interfere
15 with the activity of the active compounds therein.

 A pharmaceutically-acceptable salt form of the heterologous fusion proteins of the present invention may be used in the present invention. Acids commonly employed to form acid addition salts are inorganic acids such as
20 hydrochloric acid, hydrobromic acid, hydriodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as p-toluenesulfonic acid, methanesulfonic acid, oxalic acid, p-bromophenyl-sulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like.
25 Preferred acid addition salts are those formed with mineral acids such as hydrochloric acid and hydrobromic acid.

 Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like. Such
30 bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, and the like.

 Administration may be via any route known to be effective
35 by the physician of ordinary skill. Peripheral parenteral is one such method. Parenteral administration is commonly

understood in the medical literature as the injection of a dosage form into the body by a sterile syringe or some other mechanical device such as an infusion pump. Peripheral parenteral routes can include intravenous, intramuscular, subcutaneous, and intraperitoneal routes of administration.

The heterologous fusion proteins of the present invention may also be amenable to administration by oral, rectal, nasal, or lower respiratory routes, which are non-parenteral routes. Of these non-parenteral routes, the lower respiratory route and the oral route are preferred.

The heterologous fusion proteins of the present invention can be used to treat patients with insufficient circulating neutrophil levels, typically those undergoing cancer chemotherapy.

An "effective amount" of the heterologous fusion protein is the quantity which results in a desired therapeutic and/or prophylactic effect without causing unacceptable side-effects when administered to a subject in need of G-CSF receptor stimulation. A "desired therapeutic effect" includes one or more of the following: 1) an amelioration of the symptom(s) associated with the disease or condition; 2) a delay in the onset of symptoms associated with the disease or condition; 3) increased longevity compared with the absence of the treatment; and 4) greater quality of life compared with the absence of the treatment.

The present invention comprises G-CSF compounds that have improved biochemical and biophysical properties by virtue of being fused to an albumin protein, an albumin fragment, an albumin analog, a Fc protein, a Fc fragment, or a Fc analog. These heterologous proteins can be successfully expressed in host cells, retain signaling activities associated with activation of the G-CSF receptor, and have prolonged half-lives.

EXAMPLES**Example 1: Construction of DNA encoding heterologous fusion proteins:**

Table 1 provides the sequence of primers used to create
5 functional glycosylation sites in different regions of the
G-CSF protein (See Figure 1).

Table 1: Primer sequences used to introduce mutations into
human G-CSF.

Mutation	A Primer*	B Primer*	C Primer*	D Primer*
WT	CF177[SEQ ID NO:26] GTAAGCTTGCGT CGACGCTAGCGG CGCGCCGCCATG GCCGGACCTGCC ACCCAGAGCCCC ATGAAGCTG	CF178[SEQ ID NO:27] GGGGCAGGGAGC TGGCTGGGCCCA GTGGAGTGGCTT CCTGCACTGTCC AGAGTGCACGTG G	CF179[SEQ ID NO:28] GGACAGTGCAGG AAGCCACTCCAC TGGGCCAGCCA GCTCCCTGCCCC AGAGCTTCCTG	CF176[SEQ ID NO:29] GAACCTCGAGGA TCCTCATTAGGG CTGGGCAAGGTG CCTTAAGACGCG GTACGACACCTC CAGGAAGCTCTG
C17A SacI	CF177[SEQ ID NO:30] GTAAGCTTGCGT CGACGCTAGCGG CGCGCCGCCATG GCCGGACCTGCC ACCCAGAGCCCC ATGAAGCTG	C17Arev[SEQ ID NO:31] GCTCTAAGGCCT TGAGCAGGAAGC TCTGGGGCAGGG AGCTCGCTGGGC CCAGTGGAG	C17Afor[SEQ ID NO:32] GGGCCCAGCGAG CTCCCTGCCCCA GAGCTTCCTGCT CAAGGCCTTAGA GCAAG	CF176[SEQ ID NO:33] GAACCTCGAGGA TCCTCATTAGGG CTGGGCAAGGTG CCTTAAGACGCG GTACGACACCTC CAGGAAGCTCTG
A37N,Y39T SpeI	CF177[SEQ ID NO:34] GTAAGCTTGCGT CGACGCTAGCGG CGCGCCGCCATG GCCGGACCTGCC ACCCAGAGCCCC ATGAAGCTG	A37Nrev[SEQ ID NO:35] GTCCGAGCAGCA CTAGTTCCTCGG GGTGGCACAGCT TGGTGGTGTAC ACAGCTTCTCCT G	A37Nfor[SEQ ID NO:36] GGCGCAGCGCTC CAGGAGAAGCTG TGTAACACCACC AAGCTGTGCCAC CCCGAGGAACTA GTGCTG	CF176[SEQ ID NO:37] GAACCTCGAGGA TCCTCATTAGGG CTGGGCAAGGTG CCTTAAGACGCG GTACGACACCTC CAGGAAGCTCTG
T133N, G135T	CF177[SEQ ID NO:38]	T133Nrev[SEQ ID NO:39]	T133Nfor[SEQ ID NO:40]	CF176[SEQ ID NO:41]

<i>Eco47III</i>	GTAAGCTTGCGT CGACGCTAGCGG CGCGCCGCCATG GCCGGACCTGCC ACCCAGAGCCCC ATGAAGCTG	GCCCGGCGCTGG AAAGCGCTGGCG AAGGCCGGCATG GCGGTCTGGTTG GGCTGCAGGGCA G	GGCCCCTGCCCT GCAGCCCAACCA GACCGCCATGCC GGCCTTCGCCAG CGCTTTCCAGCG	GAACCTCGAGGA TCCTCATTAGGG CTGGGCAAGGTG CCTTAAGACGCG GTACGACACCTC CAGGAAGCTCTG
A141N, A143T <i>SapI</i>	CF177 [SEQ ID NO:42] GTAAGCTTGCGT CGACGCTAGCGG CGCGCCGCCATG GCCGGACCTGCC ACCCAGAGCCCC ATGAAGCTG	A141Nrev [SEQ ID NO:43] GCCCGGCGCTGG AAGGTAGAGTTG AAGGCCGGCATG GCACCCTGGGTG GGCTGAAGAGCA GGGGCCAT	A141Nfor [SEQ ID NO:44] GGGAATGGCCCC TGCTCTTCAGCC CAGCCAGGGTGC CATGCCGGCCTT CAACTCTACCTT CCAGCGCCGGGC AG	CF176 [SEQ ID NO:45] GAACCTCGAGGA TCCTCATTAGGG CTGGGCAAGGTG CCTTAAGACGCG GTACGACACCTC CAGGAAGCTCTG

P57V, W58N, P60T <i>HpaI</i>	JCB128 [SEQ ID NO:46] GCTAGCGGCGCG CCACCATG	JCB136 [SEQ ID NO:47] GCTCAGGGTAGC GTTAACGATGCC CAGAGAGTG	JCB137 [SEQ ID NO:48] GGGCATCGTTAA CGCTACCCTGAG CAGCTG	JCB129 [SEQ ID NO:49] GACTCGAGGATC CTCATTAGGGCT GGG
Q67N, L69T <i>NaeI</i>	JCB134 [SEQ ID NO:50] GCTAGCGGCGCG CCACCATGGCCG GACCTGCCACCC AG	JCB138 [SEQ ID NO:51] CAAGCAGCCGGC CAGCTGGGTGGC GTTGCTGGGGCA GCTGCTCAG	JCB139 [SEQ ID NO:52] GCCCCAGCAACG CCACCCAGCTGG CCGGCTGCTTGA G	JCB135 [SEQ ID NO:53] GACTCGAGGATC CTCATTAGGGCT GGGCAAGGTGCC TTAAGACGCGG
P60N, S62T <i>SpeI</i>	JCB128 [SEQ ID NO:54] GCTAGCGGCGCG CCACCATG	JCB130 [SEQ ID NO:55] GGGGCAACTAGT CAGGTTAGCCCA GGG	JCB131 [SEQ ID NO:56] GCTAACCTGACT AGTTGCCCCAGC CAG	JCB129 [SEQ ID NO:57] GACTCGAGGATC CTCATTAGGGCT GGG
S63N, P65T <i>MfeI</i>	JCB128 [SEQ ID NO:58] GCTAGCGGCGCG CCACCATG	JCB132 [SEQ ID NO:59] GGTGCAATTGCT CAGGGGAGCCCA	JCB133 [SEQ ID NO:60] GCAATTGCACCA GCCAGGCCCTG	JCB129 [SEQ ID NO:61] GACTCGAGGATC CTCATTAGGGCT

		G		GGG
E93N, I95T <i>BspEI</i>	JCB134 [SEQ ID NO:62] GCTAGCGGCGCG CCACCATGGCCG GACCTGCCACCC AG	JCB140 [SEQ ID NO:63] CCGGACTGGTCC CGTTCAGGGCCT GCAGGAGCCCCT G	JCB141 [SEQ ID NO:64] GAACGGGACCAG TCCGGAGTTGGG TCCCACCTTGG	JCB135 [SEQ ID NO:65] GACTCGAGGATC CTCATTAGGGCT GGGCAAGGTGCC TTAAGACGCGG
<i>Sall</i>	JCB155 [SEQ ID NO:66] GTCGACGCTAGC GGCGCGCCACCA TGGCCGGACCTG			

*Nucleotides in bold represent changes imposed in the target sequence and nucleotides in bold and italics represent flanking sequences which may add restriction sites to facilitate cloning, Kozac sequences, or stop codons.

Preparation 1a: DNA encoding wild-type human G-CSF

A strand overlapping extension PCR reaction was used to create a wild type human G-CSF construct in order to eliminate the methylation of an *ApaI* site. Isolated human G-CSF cDNA served as the template for these reactions. The 5' end A primer was used to create a restriction enzyme site prior to the start of the coding region as well as to introduce a Kozac sequence (GGCGCC) 5' of the coding leader sequence to facilitate translation in cell culture.

The A-B product was generated using primers CF177 and CF178 in a PCR reaction. Likewise, the C-D product was produced with primers CF179 and CF176. The products were isolated and combined. The combined mixture was then used as a template with primers CF177 and CF178 to create the full-length wild-type construct. [Nelson, R.M. and Long, G.C. (1989), *Anal. Biochem.* 180:147-151].

The full-length product was ligated into the pCR2.1-Topo vector (Invitrogen, Inc. Cat. No. K4500-40) by way of a topoisomerase TA overhang system to create pCR2.1G-CSF.

The following protocol was used for preparation of the full-length wild-type G-CSF protein as well as each of the G-CSF analogs. Approximately 5 ng of template DNA and 15 pmol of each primer was used in the initial PCR reactions.

5 The reactions were prepared using Platinum PCR Supermix® (GibcoBRL Cat. No. 11306-016). The PCR reactions were denatured at 94°C for 5 min and then subject to 25 cycles wherein each cycle consisted of 30 seconds at 94°C followed by 30 seconds at 60°C followed by 30 seconds at 72°C. A
10 final extension was carried out for 7 minutes at 72°C. PCR fragments were isolated from agarose gels and purified using a Qiaquick® gel extraction kit (Qiagen, Cat. No. #28706). DNA was resuspended in sterile water and used for the final PCR reaction to prepare full-length product.

15 Preparation 1b: DNA encoding G-CSF[A37N,Y39T,P57V,W58N,P60T,Q67N,L69T] was constructed as follows:

DNA encoding G-CSF[A37N,Y39T,Q67N,L69T] was subcloned into pJB02 to create pJB02G-CSF[A37N,Y39T,Q67N,L69T] and
20 pJB02G-CSF[A37N,Y39T,P57V,W58N,P60T] served as the template for strand overlapping expression PCR. JCB155 and JCB136 served as the A and B primers and JCB137 and JCB135 served as the C and D primers. The full-length mutated cDNA was prepared as described previously using JCB155 and JCB134
25 primers. The resulting full-length DNA encodes a protein with consensus N-linked glycosylation sites in region 1, region 2, and region 9 of the protein (See Figure 1). The full-length cDNA was ligated back into pCR2.1-Topo to create pCR2.1G-CSF[A37N,Y39T,P57V,W58N,P60T,
30 Q67N,L69T].

Preparation 1c: DNA encoding G-CSF[A37N,Y39T,S63N,P64T,E93N,I95T] was constructed as follows:

DNA encoding G-CSF[A37N,Y39T,E93N,I95T] was subcloned
35 into pJB02 to create pJB02G-CSF[A37N,Y39T,E93N,I95T] and pJB02G-CSF[A37N,Y39T,E93N,I95T] served as the template for

strand overlapping expression PCR. JCB155 and JCB132 served as the A and B primers and JCB133 and JCB135 served as the C and D primers. The full-length mutated cDNA was prepared as described previously using JCB155 and JCB135 primers. The resulting full-length DNA encodes a protein with consensus N-linked glycosylation sites in region 1, region 7, and region 10 of the protein (See Figure 1). The full-length cDNA was ligated back into pCR2.1-Topo to create pCR2.1G-CSF[A37N,Y39T,S63N,P64T,E93N,I95T].

Preparation 1d: DNA encoding G-CSF[C17A] which is G-CSF wherein the amino acid at position 17 is substituted with Ala is constructed as follows:

The wild-type construct in the pCR2.1-Topo vector (pCR2.1G-CSF) serves as the PCR template for the C17A mutagenesis. Strand overlapping extension PCR is performed as described previously. CF177 and C17Arev serve as the A-B primers and C17Afor and CF176 serve as the C-D primers. The full-length mutated cDNA is prepared as described previously using the CF177 and CF176 primers. The B and C primers are used to mutate the DNA such that a *SacI* restriction site is created and the protein expressed from the full-length sequence contains an Alanine instead of a Cysteine at position 17. The full-length cDNA is ligated back into the pCR2.1-Topo vector to create pCR2.1G-CSF[C17A] wherein the sequence is confirmed. G-CSF analog encoding DNA is then cloned into the *Nhe/Xho* sites of mammalian expression vector pJB02 to create pJB02G-CSF[C17A].

Preparation 1e: DNA encoding G-CSF[A37N,Y39T] is constructed as follows:

Strand overlapping extension PCR is performed using pCR2.1G-CSF[C17A] as the template. Primers CF177 and A37Nrev serve as the A-B primers and CF176 and A37Nfor serve as the C-D primers. The full-length mutated cDNA is prepared as described previously using the CF177 and CF176 primers.

The B and C primers contain mismatched sequences such that a *SpeI* site is created in the DNA and the protein expressed from the full-length sequence contains a consensus sequence for N-linked glycosylation in region 1 of the protein. The full-length cDNA is ligated back into the pCR2.1-Topo vector to create pCR2.1G-CSF[A37N,Y39T] wherein the sequence is confirmed. G-CSF analog encoding DNA is then cloned into the *Nhe/Xho* sites of mammalian expression vector pJB02 to create pJB02G-CSF[A37N,Y39T].

Preparation 1f: DNA encoding G-CSF[P57V,W58N,P60T] is constructed as follows:

Strand overlapping extension PCR is performed using pJB02G-CSF[C17A] as the template. Primers JCB128 and JCB136 serve as the A-B primers and JCB137 and JCB129 serve as the C-D primers. The full-length mutated cDNA is prepared as described previously using the JCB128 and JCB129 primers. The B and C primers contain mismatched sequences such that a *HpaI* site is created and the protein expressed from the full-length sequence contains a consensus sequence for N-linked glycosylation in region 2 of the protein. The full-length cDNA is ligated back into the pCR2.1-Topo vector to create pCR2.1G-CSF[P57V,W58N,P60T] wherein the sequence is confirmed. G-CSF analog encoding DNA is then cloned into the *Nhe/Xho* sites of mammalian expression vector pJB02 to create pJB02G-CSF[P57V,W58N,P60T].

Preparation 1g: DNA encoding G-CSF[P60N,S62T] is constructed as follows:

Strand overlapping extension PCR is performed using pJB02G-CSF[C17A] as the template. Primers JCB128 and JCB130 serve as the A-B primers and JCB131 and JCB129 serve as the C-D primers. The full-length mutated cDNA is prepared as described previously using the JCB128 and JCB129 primers. The B and C primers contain mismatched sequences such that a *SpeI* site is created and the protein expressed from the

full-length sequence contains a consensus sequence for N-linked glycosylation in region 4 of the protein. The full-length cDNA is ligated back into the pCR2.1-Topo vector to create pCR2.1G-CSF[P60N,S62T] wherein the sequence is confirmed. G-CSF analog encoding DNA is then cloned into the *Nhe/Xho* sites of mammalian expression vector pJB02 to create pJB02G-CSF[P60N,S62T].

Preparation 1h: DNA encoding G-CSF[S63N,P65T] is constructed as follows:

Strand overlapping extension PCR is performed using pJB02G-CSF[C17A] as the template. Primers JCB128 and JCB132 serve as the A-B primers and JCB133 and JCB129 serve as the C-D primers. The full-length mutated cDNA is prepared as described previously using the JCB128 and JCB129 primers. The B and C primers contain mismatched sequences such that a *MfeI* site is created and the protein expressed from the full-length sequence contains a consensus sequence for N-linked glycosylation in region 7 of the protein. The full-length cDNA is ligated back into the pCR2.1-Topo vector to create pCR2.1G-CSF[S63N,P65T] wherein the sequence is confirmed. G-CSF analog encoding DNA is then cloned into the *Nhe/Xho* sites of mammalian expression vector pJB02 to create pJB02G-CSF[S63N,P65T].

Preparation 1i: DNA encoding G-CSF[Q67N,L69T] is constructed as follows:

Strand overlapping extension PCR is performed using pJB02G-CSF[C17A] as the template. Primers JCB134 and JCB138 serve as the A-B primers and JCB139 and JCB135 serve as the C-D primers. The full-length mutated cDNA is prepared as described previously using the JCB128 and JCB129 primers. The B and C primers contain mismatched sequences such that a *NaeI* site is created and the protein expressed from the full-length sequence contains a consensus sequence for N-linked glycosylation in region 9 of the protein. The full-

length cDNA is ligated back into the pCR2.1-Topo vector to create pCR2.1G-CSF[Q67N,L69T] wherein the sequence is confirmed. G-CSF analog encoding DNA is then cloned into the *Nhe/Xho* sites of mammalian expression vector pJB02 to create
5 pJB02G-CSF[Q67N,L69T].

Preparation 1j: DNA encoding G-CSF[E93N,I95T] is constructed as follows:

Strand overlapping extension PCR is performed using
10 pJB02G-CSF[C17A] as the template. Primers JCB134 and JCB140 serve as the A-B primers and JCB141 and JCB135 serve as the C-D primers. The full-length mutated cDNA is prepared as described previously using the JCB128 and JCB129 primers. The B and C primers contain mismatched sequences such that
15 a *BspEI* site is created and the protein expressed from the full-length sequence contains a consensus sequence for N-linked glycosylation in region 10 of the protein. The full-length cDNA is ligated back into the pCR2.1-Topo vector to create pCR2.1G-CSF[E93N,I95T] wherein the sequence is
20 confirmed. G-CSF analog encoding DNA is then cloned into the *Nhe/Xho* sites of mammalian expression vector pJB02 to create pJB02G-CSF[E93T,I95T].

Preparation 1k: DNA encoding G-CSF[T133N,G135T] is
25 constructed as follows:

Strand overlapping extension PCR is performed using pCR2.1G-CSF[C17A] as the template. Primers CF177 and T133Nrev serve as the A-B primers and T133Nfor and CF176 serve as the C-D primers. The full-length mutated cDNA is
30 prepared as described previously using the CF177 and CF176 primers. The B and C primers contain mismatched sequences such that an *Eco47III* site is created and the protein expressed from the full-length sequence contains a consensus sequence for N-linked glycosylation in region 13 of the
35 protein. The full-length cDNA is ligated back into the

pCR2.1-Topo vector to create pCR2.1G-CSF[T133N,G135T] wherein the sequence is confirmed.

Preparation 1l: DNA encoding G-CSF[A141N,A143T] is
5 constructed as follows:

Strand overlapping extension PCR is performed using pCR2.1G-CSF[C17A] as the template. Primers CF177 and A141Nrev serve as the A-B primers and A141Nfor and CF176 serve as the C-D primers. The full-length mutated cDNA is
10 prepared as described previously using the CF177 and CF176 primers. The B and C primers contain mismatched sequences such that an *SapI* site is created and the protein expressed from the full-length sequence contains a consensus sequence for N-linked glycosylation in region 14 of the protein. The
15 full-length cDNA is ligated back into the pCR2.1-Topo vector to create pCR2.1G-CSF[A141N,A143T] wherein the sequence is confirmed.

Preparation 1m: DNA encoding G-CSF[A37N,Y39T,T133N,
20 G135T] is constructed as follows:

A 210 bp insert containing G-CSF[A37N,Y39T] is isolated from pCR2.1G-CSF[A37N,Y39T] using *EcoNI*. This fragment is ligated into pCR2.1G-CSF[T133N,G135T] which is prepared by cleavage with *EcoNI* and subsequent isolation of the vector
25 (4359 bp) from a 210 bp fragment containing wild-type G-CSF sequences. This ligation creates pCR2.1G-CSF[A37N,Y39T,T133N,G135T]. Analog encoding DNA is then subcloned into pJB02 using *NheI/XhoI* to create pJB02G-CSF[A37N,Y39T,T133N,G135T].

30

Preparation 1n: DNA encoding G-CSF[A37N,Y39T,A141N,A143T] is constructed as follows:

A 210 bp insert containing G-CSF[A37N,Y39T] is isolated from pCR2.1G-CSF[A37N,Y39T] using *EcoNI*. This fragment is
35 ligated into pCR2.1G-CSF[A141N,A143T] which is prepared by cleavage with *EcoNI* and subsequent isolation of the vector

(4359 bp) from a 210 bp fragment containing wild-type G-CSF sequences. This ligation creates pCR2.1G-CSF[A37N,Y39T,A141N,A143T]. Analog encoding DNA is then subcloned into pJB02 using *NheI/XhoI* to create pJB02G-CSF[A37N,Y39T,A141N,A143T].

Preparation 1o: DNA encoding G-CSF[A37N,Y39T,P57V,W58N,P60T] is constructed as follows:

DNA encoding G-CSF[A37N,Y39T] is subcloned into pJB02 to create pJB02G-CSF[A37N,Y39T] and pJB02G-CSF[A37N,Y39T] serves as the template for strand overlapping expression PCR. JCB128 and JCB136 serve as the A and B primers and JCB137 and JCB129 serve as the C and D primers. The full-length mutated cDNA is prepared as described previously using JCB128 and JCB129 primers. The resulting full-length DNA encodes a protein with consensus N-linked glycosylation sites in region 1 and region 2 of the protein. The full-length cDNA is ligated back into pCR2.1-Topo to create pCR2.1G-CSF[A37N,Y39T,P57V,W58N,P60T].

Preparation 1p: DNA encoding G-CSF[A37N,Y39T,Q67N,L69T] is constructed as follows:

DNA encoding G-CSF[A37N,Y39T] is subcloned into pJB02 to create pJB02G-CSF[A37N,Y39T] and pJB02G-CSF[A37N,Y39T] serves as the template for strand overlapping expression PCR. JCB134 and JCB138 serve as the A and B primers and JCB139 and JCB135 serve as the C and D primers. The full-length mutated cDNA is prepared as described previously using JCB128 and JCB129 primers. The resulting full-length DNA encodes a protein with consensus N-linked glycosylation sites in region 1 and region 9 of the protein. The full-length cDNA is ligated back into pCR2.1-Topo to create pCR2.1G-CSF[A37N,Y39T,Q67N,L69T].

Preparation 1q: DNA encoding G-CSF[A37N,Y39T,E93N,I95T] is constructed as follows:

DNA encoding G-CSF[A37N,Y39T] is subcloned into pJB02 to create pJB02G-CSF[A37N,Y39T] and pJB02G-CSF[A37N,Y39T] serves as the template for strand overlapping expression PCR. JCB134 and JCB140 serve as the A and B primers and JCB141 and JCB135 serve as the C and D primers. The full-length mutated cDNA is prepared as described previously using JCB128 and JCB129 primers. The resulting full-length DNA encodes a protein with consensus N-linked glycosylation sites in region 1 and region 10 of the protein. The full-length cDNA is ligated back into pCR2.1-Topo to create pCR2.1G-CSF[A37N,Y39T,E93N,I95T].

Various mutated G-CSF polypeptides were then ligated in-frame directly to DNA encoding IgG1 or IgG4. Various mutated G-CSF polypeptides were also ligated in-frame via DNA encoding a peptide linker or directly to human albumin.

Example 2: Expression of heterologus fusion proteins:

2a: Expression in 293/EBNA cells:

Each full-length DNA encoding a heterologous fusion protein was cloned into the pJB02. This vector contains both the Ori P and Epstein Barr virus nuclear antigen (EBNA) components which are necessary for sustained, transient expression in 293 EBNA cells. Further, this expression plasmid contains a puromycin resistance gene expressed from the CMV promoter as well as an ampicillin resistance gene. The gene of interest is also expressed from the CMV promoter.

The transfection mixture was prepared by mixing 73 μ l of the liposome transfection agent Fugene 6® (Roche Molecular Biochemicals, Cat. No. 1815-075) with 820 μ l Opti-Mem® (GibcoBRL Cat. No. 31985-062). G-CSF pJB02 DNA (12 μ g), prepared using a Qiagen plasmid maxiprep kit (Qiagen, Cat. No. 12163), was then added to the mixture.

The mixture was incubated at room temperature for 15 minutes.

Cells were plated on 10 cm² plates in DMEM/F12 3:1 (GibcoBRL Cat. No. 93-0152DK) supplemented with 5% fetal bovine serum, 20mM HEPES, 2 mM L-glutamine, and 50 µg/mL Geneticin such that the plates were 60% to 80% confluent by the time of the transfection. Immediately before the transfection mixture was added to the plates, fresh media was added. The mixture was then added dropwise to cells with intermittent swirling. Plates were then incubated at 37°C in a 5% CO₂ atmosphere for 24 hours at which point the media was changed to hybritech medium without serum. The media containing a secreted form of a glycosylated G-CSF analog was then isolated 48 hours later.

2b: Expression in CHO cells:

The expression vector for expression in CHO-K1 cells is pEE14.1. This vector includes the glutamine synthetase gene which enables selection using methionine sulfoximine. This gene includes two poly A signals at the 3' end. G-CSF analogs are expressed from the CMV promoter which includes 5' untranslated sequences from the hCMV-MIE gene to enhance mRNA levels and translatability. The SV40 poly A signal is cloned 3' of the fusion protein DNA. The SV40 late promoter drives expression of GS minigene. This expression vector encoding the gene of interest is prepared for transfection using a QIAGEN Maxi Prep Kit (QIAGEN, Cat. No. 12362). The final DNA pellet (50-100 µg) is resuspended in 100 µl of basal formulation medium (GibcoBRL CD-CHO Medium without L-Glutamine, without thymidine, without hypoxanthine).

Before each transfection, CHO-K1 cells are counted and checked for viability. A volume equal to 1 x 10⁷ cells is centrifuged and the cell pellet rinsed with basal formulation medium. The cells are centrifuged a second time and the final pellet resuspended in basal formulation medium (700 µl final volume).

The resuspended DNA and cells are then mixed together in a standard electroporation cuvette (Gene Pulsar Cuvette) used to support mammalian transfections, and placed on ice for five minutes. The cell/DNA mix is then electroporated in a BioRad Gene Pulsar device set at 300V/975 μ F and the cuvette placed back on ice for five minutes. The cell/DNA mixed is diluted into 20 ml of cell growth medium in a non-tissue culture treated T75 flask and incubated at 37°C / 5% CO₂ for 48-72 hours.

The cells are counted, checked for viability, and plated at various cell densities in selective medium in 96 well tissue culture plates and incubated at 37°C in a 5% CO₂ atmosphere. Selective medium is basal medium with 1X HT Supplement (GibcoBRL 100X HT Stock), 100 μ g/mL Dextran Sulfate (Sigma 100 mg/ml stock), 1X GS Supplements (JRH BioSciences 50X Stock) and 25 μ M MSX (Methionine Sulphoximine). The plates are monitored for colony formation and screened for glycosylated G-CSF analog production.

Example 3: Purification of Heterologous Fusion Proteins

Heterologous fusion proteins containing a hyperglycosylated G-CSF protein fused to albumin were harvested from host cells and then dialyzed against 20 mM Tris pH 7.4. An anion exchange column (1 ml Pharmacia HiTrap Q) was equilibrated with 20 mM Tris pH 7.4 and the dialyzed material loaded at 2 ml/min. The protein was eluted from the column using a linear gradient from 0 to 500 mM NaCl in 80 min at 1 ml/min and elution was monitored by UV absorbance at 280 nm. SDS-PAGE analysis was used to identify and pool fractions of interest. This pool was dialyzed against 25 mM sodium acetate (NaOAc) pH 5.0

A cation exchange column (1 ml Pharmacia HiTrap S column) was equilibrated with 25 mM NaOAc pH 5.0 and the dialysate was loaded at 1 ml/min. The protein was eluted

from the column using a linear gradient from 0 to 500 mM NaCl in 30 min. The fractions were immediately neutralized with 1 M Tris pH 8 to a final pH of 7. SDS-PAGE gels were used to identify and pool fractions of interest.

- 5 Heterologous fusion proteins containing a hyperglycosylated G-CSF protein fused to an Fc protein were harvested from host cells and dialyzed against 20 mM sodium phosphate pH 7.0. An affinity column (1 ml Pharmacia HiTrap Protein A or rProtein A) was equilibrated with 20 mM sodium phosphate pH 7.0 and the dialysate was loaded at 2 ml/min. 10 1 ml/min of 100 mM citric acid pH 3 was used to elute the protein. Fractions were immediately neutralized with 1M Tris pH 8 to pH 7 and peak fractions (determined by in-line OD280 monitoring) were further diluted with 20 mM sodium 15 phosphate pH 7.0. . SDS-PAGE analysis was used to identify and pool fractions of interest.

235323 "2346789"

WE CLAIM:

1. A heterologous fusion protein comprising a
 5 hyperglycosylated G-CSF analog fused to a polypeptide
 selected from the group consisting of
 - a) human albumin;
 - b) human albumin analogs; and
 - c) fragments of human albumin.
- 10 2. The heterologous fusion protein of claim 1, wherein the
 hyperglycosylated G-CSF analog is fused to the polypeptide
 via a peptide linker.
- 15 3. The heterologous fusion protein of Claim 2 wherein the
 peptide linker is selected from the group consisting of:
 - a) a glycine rich peptide;
 - b) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]_n
 where n is 1, 2, 3, 4, or 5; and
 - 20 c) a peptide having the sequence [Gly-Gly-Gly-Gly-
 Ser]₃.
4. The heterologous fusion protein of Claims 1, 2, or 3
 wherein the hyperglycosylated G-CSF analog comprises the
 25 amino acid sequence of the formula I: [SEQ ID NO: 1]

1	5	10	15													
Thr	Pro	Leu	Gly	Pro	Ala	Ser	Ser	Leu	Pro	Gln	Ser	Phe	Leu	Leu	Lys	
			20					25					30			
30	Xaa	Leu	Glu	Gln	Val	Arg	Lys	Ile	Gln	Gly	Asp	Gly	Ala	Ala	Leu	Gln
			35					40					45			
	Glu	Lys	Leu	Cys	Xaa	Xaa	Xaa	Lys	Leu	Cys	His	Pro	Glu	Glu	Leu	Val
			50					55				60				
	Leu	Leu	Gly	His	Ser	Leu	Gly	Ile	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
35			65				70				75				80	
	Xaa	Xaa	Xaa	Xaa	Xaa	Gln	Leu	Ala	Gly	Cys	Leu	Ser	Gln	Leu	His	Ser
					85				90				95			
	Gly	Leu	Phe	Leu	Tyr	Gln	Gly	Leu	Leu	Gln	Ala	Leu	Xaa	Xaa	Xaa	Ser
			100					105					110			
40	Xaa	Glu	Leu	Gly	Pro	Thr	Leu	Asp	Thr	Leu	Gln	Leu	Asp	Val	Ala	Asp
			115					120					125			
	Phe	Ala	Thr	Thr	Ile	Trp	Gln	Gln	Met	Glu	Glu	Leu	Gly	Met	Ala	Pro
			130				135					140				
	Ala	Leu	Gln	Pro	Xaa	Xaa	Xaa	Ala	Met	Pro	Ala	Phe	Xaa	Xaa	Xaa	Phe
45			145				150				155					160

Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser His Leu Aln Ser Phe
 165 170
 Leu Glu Val Ser Tyr Arg Val Leu Arg His Leu Ala Gln Pro (I)

5 wherein:

Xaa at position 17 is Cys, Ala, Leu, Ser, or Glu;
 Xaa at position 37 is Ala or Asn;
 Xaa at position 38 is Thr, or any other amino acid except
 Pro;

10 Xaa at position 39 is Tyr, Thr, or Ser;

Xaa at position 57 is Pro or Val;

Xaa at position 58 is Trp or Asn;

Xaa at position 59 is Ala or any other amino acid except
 Pro;

15 Xaa at position 60 is Pro, Thr, Asn, or Ser,

Xaa at position 61 is Leu, or any other amino acid except
 Pro;

Xaa at position 62 is Ser or Thr;

Xaa at position 63 is Ser or Asn;

20 Xaa at position 64 is Cys or any other amino acid except
 Pro;

Xaa at position 65 is Pro, Ser, or Thr;

Xaa at position 66 is Ser or Thr;

Xaa at position 67 is Gln or Asn;

25 Xaa at position 68 is Ala or any other amino acid except
 Pro;

Xaa at position 69 is Leu, Thr, or Ser

Xaa at position 93 is Glu or Asn

Xaa at position 94 is Gly or any other amino acid except

30 Pro;

Xaa at position 95 is Ile, Asn, Ser, or Thr;

Xaa at position 97 is Pro, Ser, Thr, or Asn;

Xaa at position 133 is Thr or Asn;

Xaa at position 134 is Gln or any other amino acid except

35 Pro;

Xaa at position 135 is Gly, Ser, or Thr

Xaa at position 141 is Ala or Asn;

Xaa at position 142 is Ser or any other amino acid except
 Pro; and

40 Xaa at position 143 is Ala, Ser, or Thr;

and wherein:

Xaa at positions 37, 38, and 39 constitute region 1;

Xaa at positions 58, 59, and 60 constitute region 2;

45 Xaa at positions 59, 60, and 61 constitute region 3;

Xaa at positions 60, 61, and 62 constitute region 4;

- 5 Xaa at positions 61, 62, and 63 constitute region 5;
Xaa at positions 62, 63, and 64 constitute region 6;
Xaa at positions 63, 64, and 65 constitute region 7;
Xaa at positions 64, 65, and 66 constitute region 8;
Xaa at positions 67, 68, and 69 constitute region 9;
Xaa at positions 93, 94, and 95 constitute region 10;
Xaa at positions 94, 95, and Ser at position 96
constitute region 11;
Xaa at positions 95, and 97, and Ser at position 96
10 constitute region 12;
Xaa at positions 133, 134, and 135 constitute
region 13;
Xaa at positions 141, 142, and 143 constitute
region 14;

15 and provided that at least one of regions 1 through 14
comprises the sequence Asn Xaa1 Xaa2 wherein Xaa1 is any
amino acid except Pro and Xaa2 is Ser or Thr.

- 20 5. The heterologous fusion protein of Claim 4 wherein any
two regions of regions 1 through 14 comprise the sequence
Asn Xaa1 Xaa2 wherein Xaa1 is any amino acid except Pro and
Xaa2 is Ser or Thr.
- 25 6. The heterologous fusion protein of Claim 4 wherein any
three regions of regions 1 through 14 comprise the sequence
Asn Xaa1 Xaa2 wherein Xaa1 is any amino acid except Pro and
Xaa2 is Ser or Thr.
- 30 7. The heterologous fusion protein of Claim 4 wherein any
four regions of regions 1 through 14 comprise the sequence
Asn Xaa1 Xaa2 wherein Xaa1 is any amino acid except Pro and
Xaa2 is Ser or Thr

8. The heterologous fusion protein of Claim 4 wherein the hyperglycosylated G-CSF analog is selected from the group consisting of:

- a) G-CSF[A37N,Y39T]
- 5 b) G-CSF[P57V,W58N,P60T]
- c) G-CSF[P60N,S62T]
- d) G-CSF[S63N,P65T]
- e) G-CSF[Q67N,L69T]
- f) G-CSF[E93N,I95T]
- 10 g) G-CSF[T133N,G135T]
- h) G-CSF[A141N,A143T]
- i) G-CSF[A37N,Y39T,P57V,W58N,P60T]
- j) G-CSF[A37N,Y39T,P60N,S62T]
- k) G-CSF[A37N,Y39T,S63N,P65T]
- 15 l) G-CSF[A37N,Y39T,Q67N,L69T]
- m) G-CSF[A37N,Y39T,E93N,I95T]
- n) G-CSF[A37N,Y39T,T133N,G135T]
- o) G-CSF[A37N,Y39T,A141N,A143T]
- p) G-CSF[A37N,Y39T,P57V,W58N,P60T,S63N,P65T]
- 20 q) G-CSF[A37N,Y39T,P57V,W58N,P60T,Q67N,L69T]
- r) G-CSF[A37N,Y39T,S63N,P65T,E93N,I95T]

9. The heterologous fusion protein of claim 8, wherein the hyperglycosylated G-CSF analog is G-CSF[A37N,
25 Y39T,P57V,W58N,P60T,Q67N,L69T].

10. The heterologous fusion protein of claim 8, wherein the hyperglycosylated G-CSF analog is G-
30 CSF[A37N,Y39T,S63N,P65T,E93N,I95T].

11. A heterologous fusion protein which is the product of the expression in a host cell of an exogenous DNA sequence which comprises a DNA sequence encoding a heterologous
35 fusion protein of any one of Claims 1 through 10.

12. A polynucleotide encoding a heterologous fusion protein of any one of Claims 1 through 11.

13. A polynucleotide which comprises a DNA sequence

5 selected from the group consisting of:

- a) SEQ ID NO:2
- b) SEQ ID NO:3
- c) SEQ ID NO:4
- d) SEQ ID NO:5
- 10 e) SEQ ID NO:6
- f) SEQ ID NO:7
- g) SEQ ID NO:8
- h) SEQ ID NO:9
- i) SEQ ID NO:10
- 15 j) SEQ ID NO:11
- k) SEQ ID NO:12
- l) SEQ ID NO:13
- m) SEQ ID NO:14
- n) SEQ ID NO:15
- 20 o) SEQ ID NO:16,

fused in-frame to a DNA encoding a protein selected from the group consisting of:

- a) human albumin,
- b) human albumin analog; and
- 25 c) fragments of human albumin.

14. The polynucleotide of Claim 13, wherein the DNA fused in-frame comprises SEQ ID NO: 17.

30 15. The heterologous fusion protein of any one of Claims 1 through 11 wherein the polypeptide is human albumin.

16. The heterologous fusion protein of any one of Claims 1 through 11 wherein the polypeptide is an N-terminal fragment
35 of albumin.

17. A method for increasing neutrophil levels in a mammal comprising administering a therapeutically effective amount of the heterologous fusion protein of any one of Claims 1
40 through 11, 15 and 16.

18. The use of the heterologous fusion protein as claimed in any one of Claims 1 through 11, 15 and 16 for the manufacture of a medicament for the treatment of patients with insufficient circulating neutrophil levels.

19. Use of a heterologous fusion protein of any one of Claims 1 through 11, 15, and 16 as a medicament.

20. Use of a heterologous fusion protein of any one of Claims 1 through 11, 15, and 16 for the treatment of patients with insufficient circulating neutrophil levels.

21. A pharmaceutical formulation adapted for the treatment of patients with insufficient neutrophil levels comprising a heterologous fusion protein of any one of Claims 1 through 11, 15, and 16.

22. A heterologous fusion protein comprising a hyperglycosylated G-CSF analog fused to a polypeptide selected from the group consisting of:

- a) the Fc portion of an immunoglobulin;
- b) an analog of the Fc portion of an immunoglobulin;
- and
- c) fragments of the Fc portion of an immunoglobulin.

23. The heterologous fusion protein of Claim 22, wherein the hyperglycosylated G-CSF analog is fused to the polypeptide via a peptide linker.

24. The heterologous fusion protein of the Claim 23 wherein the peptide linker is selected from the group consisting of:

- a) a glycine rich peptide;
- b) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]_n where n is 1, 2, 3, 4, or 5; and
- c) a peptide having the sequence [Gly-Gly-Gly-Gly-Ser]₃.

25. The heterologous fusion protein of Claims 22, 23 or 24, wherein the hyperglycosylated G-CSF analog comprises the amino acid sequence of the formula I: [SEQ ID NO: 1]

5	1	5	10	15
	Thr	Pro	Leu	Gly
		20	25	30
	Xaa	Leu	Glu	Gln
10	35	40	45	50
	Glu	Lys	Leu	Cys
	55	60	65	70
	Leu	Leu	Gly	His
	75	80	85	90
	Xaa	Xaa	Xaa	Xaa
15	95	100	105	110
	Gly	Leu	Phe	Leu
	115	120	125	130
	Xaa	Glu	Leu	Gly
20	135	140	145	150
	Phe	Ala	Thr	Thr
	155	160	165	170
	Gln	Arg	Arg	Ala
25	175	180	185	190
	Leu	Glu	Val	Ser

(I)

wherein:

- Xaa at position 17 is Cys, Ala, Leu, Ser, or Glu;
- 30 Xaa at position 37 is Ala or Asn;
- Xaa at position 38 is Thr, or any other amino acid except Pro;
- Xaa at position 39 is Tyr, Thr, or Ser;
- Xaa at position 57 is Pro or Val;
- 35 Xaa at position 58 is Trp or Asn;
- Xaa at position 59 is Ala or any other amino acid except Pro;
- Xaa at position 60 is Pro, Thr, Asn, or Ser,
- Xaa at position 61 is Leu, or any other amino acid except
- 40 Pro;
- Xaa at position 62 is Ser or Thr;
- Xaa at position 63 is Ser or Asn;
- Xaa at position 64 is Cys or any other amino acid except Pro;
- 45 Xaa at position 65 is Pro, Ser, or Thr;
- Xaa at position 66 is Ser or Thr;
- Xaa at position 67 is Gln or Asn;
- Xaa at position 68 is Ala or any other amino acid except Pro;
- 50 Xaa at position 69 is Leu, Thr, or Ser

- Xaa at position 93 is Glu or Asn
 Xaa at position 94 is Gly or any other amino acid except Pro;
 Xaa at position 95 is Ile, Asn, Ser, or Thr;
 5 Xaa at position 97 is Pro, Ser, Thr, or Asn;
 Xaa at position 133 is Thr or Asn;
 Xaa at position 134 is Gln or any other amino acid except Pro;
 Xaa at position 135 is Gly, Ser, or Thr
 10 Xaa at position 141 is Ala or Asn;
 Xaa at position 142 is Ser or any other amino acid except Pro; and
 Xaa at position 143 is Ala, Ser, or Thr;
- 15 and wherein:
 Xaa at positions 37, 38, and 39 constitute region 1;
 Xaa at positions 58, 59, and 60 constitute region 2;
 Xaa at positions 59, 60, and 61 constitute region 3;
 Xaa at positions 60, 61, and 62 constitute region 4;
 20 Xaa at positions 61, 62, and 63 constitute region 5;
 Xaa at positions 62, 63, and 64 constitute region 6;
 Xaa at positions 63, 64, and 65 constitute region 7;
 Xaa at positions 64, 65, and 66 constitute region 8;
 Xaa at positions 67, 68, and 69 constitute region 9;
 25 Xaa at positions 93, 94, and 95 constitute region 10;
 Xaa at positions 94, 95, and Ser at position 96
 constitute region 11;
 Xaa at positions 95, and 97, and Ser at position 96
 constitute region 12;
 30 Xaa at positions 133, 134, and 135 constitute
 region 13;
 Xaa at positions 141, 142, and 143 constitute
 region 14;
- 35 and provided that at least one of regions 1 through 14
 comprises the sequence Asn Xaa1 Xaa2 wherein Xaa1 is any
 amino acid except Pro and Xaa2 is Ser or Thr.

26. The heterologous fusion protein of Claim 25 wherein any two regions of regions 1 through 14 comprise the sequence Asn Xaa1 Xaa2 wherein Xaa1 is any amino acid except Pro and
5 Xaa2 is Ser or Thr.

27. The heterologous fusion protein of Claim 25 wherein any three regions of regions 1 through 14 comprise the sequence Asn Xaa1 Xaa2 wherein Xaa1 is any amino acid except Pro and
10 Xaa2 is Ser or Thr.

28. The heterologous fusion protein of Claim 25 wherein any four regions of regions 1 through 14 comprise the sequence Asn Xaa1 Xaa2 wherein Xaa1 is any amino acid except Pro and
15 Xaa2 is Ser or Thr

29. The heterologous fusion protein of Claim 25 wherein the hyperglycosylated G-CSF analog is selected from the group consisting of:

- 20 a) G-CSF[A37N,Y39T]
b) G-CSF[P57V,W58N,P60T]
c) G-CSF[P60N,S62T]
d) G-CSF[S63N,P65T]
e) G-CSF[Q67N,L69T]
25 f) G-CSF[E93N,I95T]
g) G-CSF[T133N,G135T]
h) G-CSF[A141N,A143T]
i) G-CSF[A37N,Y39T,P57V,W58N,P60T]
j) G-CSF[A37N,Y39T,P60N,S62T]
30 k) G-CSF[A37N,Y39T,S63N,P65T]
l) G-CSF[A37N,Y39T,Q67N,L69T]
m) G-CSF[A37N,Y39T,E93N,I95T]
n) G-CSF[A37N,Y39T,T133N,G135T]
o) G-CSF[A37N,Y39T,A141N,A143T]
35 p) G-CSF[A37N,Y39T,P57V,W58N,P60T,S63N,P65T]
q) G-CSF[A37N,Y39T,P57V,W58N,P60T,Q67N,L69T]

r) G-CSF[A37N,Y39T,S63N,P65T,E93N,I95T]

30. The heterologous fusion protein of Claim 29 wherein the hyperglycosylated G-CSF analog is G-

5 CSF[A37N,Y39T,P57V,W58N,P60T,Q67N,L69T]

31. The heterologous fusion protein of Claim 29 wherein the hyperglycosylated G-CSF analog is G-

CSF[A37N,Y39T,S63N,P65T,E93N,I95T]

10

32. The heterologous fusion protein of any one of Claims 22 through 31 wherein the polypeptide is the Fc portion of an Ig selected from the group consisting of: IgG1, IgG2, IgG3, IgG4, IgE, IgA, IgD, or IgM.

15

33. The heterologous fusion protein of Claim 32 wherein the polypeptide is the Fc portion of an Ig selected from the group consisting of: IgG1, IgG2, IgG3, and IgG4.

20

34. The heterologous fusion protein of Claim 33 wherein the polypeptide is the Fc portion of an IgG1 immunoglobulin.

35. The heterologous fusion protein of Claim 33 wherein the polypeptide is the Fc portion of an IgG4

25

immunoglobulin.

36. The heterologous fusion protein of any one of Claims 22 through 35 wherein the Fc portion is a human IgG protein.

30 37. The heterologous fusion protein of any one of Claims 22 through 36 wherein the Fc portion comprises hinge, CH2, and CH3 domains.

38. The heterologous fusion protein of Claim 34 wherein the
35 polypeptide has the sequence of SEQ ID NO: 33.

35

48. Use of a heterologous fusion protein of any one of Claims 22 through 29 as a medicament.

5 49. Use of a heterologous fusion protein of any one of Claims 22 through 29 for the treatment of patients with insufficient circulating neutrophil levels.

10 50. A pharmaceutical formulation adapted for the treatment of patients with insufficient neutrophil levels comprising a heterologous fusion protein of any one of Claims 22 through 29.

15 51. A heterologous fusion protein as hereinbefore described with reference to any one of the Examples.

Abstract

5 The present invention encompasses heterologous fusion
to proteins comprising a hyperglycsoylated G-CSF analog fused
immunoglobulin which act to extend the *in vivo* half-life of
the protein compared to native G-CSF. These fusion proteins
are particularly suited for the treatment of conditions
10 treatable by stimulation of circulating neutrophils, such as
after chemotherapy regimens or in chronic congenital
neutropenia.

2050000 2461303

Fig. 1

1/1

1 5 10 15
 Thr Pro Leu Gly Pro Ala Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys

 20 25 30
 Cys Leu Glu Gln Val Arg Lys Ile Gln Gly Asp Gly Ala Ala Leu Gln

 35 R1 40 45
 Glu Lys Leu Cys Xaa Xaa Xaa Lys Leu Cys His Pro Glu Glu Leu Val

 50 55 R3 60 R6 65
 Leu Leu Gly His Ser Leu Gly Ile Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Ser
 R4 R7 R8
 R2 R5 R6

 R9 70 75 80
Xaa Xaa Xaa Gln Leu Ala Gly Cys Leu Ser Gln Leu His Ser

 85 90 R12
 Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Xaa Xaa Xaa Ser Xaa
 R10 R11

 100 105 110
 Xaa Xaa Gly Pro Thr Leu Asp Thr Leu Gln Leu Asp Val Ala Asp

 115 120 125
 Phe Ala Thr Thr Ile Trp Gln Gln Met Glu Glu Leu Gly Met Ala Pro

 130 R13 135 140 R14
 Ala Leu Gln Pro Xaa Xaa Xaa Ala Met Pro Ala Phe Xaa Xaa Xaa Phe

 145 150 155 160
 Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser His Leu Aln Ser Phe

 165 170
 Leu Glu Val Ser Tyr Arg Val Leu Arg His Leu Ala Gln Pro

2050-2450
 2450-2650
 2650-2850
 2850-3050
 3050-3250
 3250-3450
 3450-3650
 3650-3850
 3850-4050
 4050-4250
 4250-4450
 4450-4650
 4650-4850
 4850-5050
 5050-5250
 5250-5450
 5450-5650
 5650-5850
 5850-6050
 6050-6250
 6250-6450
 6450-6650
 6650-6850
 6850-7050
 7050-7250
 7250-7450
 7450-7650
 7650-7850
 7850-8050
 8050-8250
 8250-8450
 8450-8650
 8650-8850
 8850-9050
 9050-9250
 9250-9450
 9450-9650
 9650-9850
 9850-10050

P-15648.ST25.txt
SEQUENCE LISTING

<110> Beals, John
Kuchibhotla, Uma

<120> HETEROLOGOUS G-CSF FUSION PROTEINS

<130> P-15648

<160> 66

<170> PatentIn version 3.1

<210> 1

<211> 174

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<220>

<221> MISC_FEATURE

<222> (17)..(17)

<223> Xaa at position 17 is Cys, Ala, Leu, Ser, or Glu;

<220>

<221> MISC_FEATURE

<222> (37)..(37)

<223> Xaa at position 37 is Ala or Asn;

<220>

<221> MISC_FEATURE

<222> (38)..(38)

<223> Xaa at position 38 is Thr, or any other amino acid exept Pro;

<220>

<221> MISC_FEATURE

<222> (39)..(39)

<223> Xaa at position 39 is Tyr, Thr, or Ser;

<220>

<221> MISC_FEATURE

<222> (57)..(57)

<223> Xaa at position 57 is Pro or Val;

<220>

<221> MISC_FEATURE

<222> (58)..(58)

<223> Xaa at position 58 is Trp or Asn;

<220>

<21> MISC_FEATURE

<222> (59)..(59)

<223> Xaa at position 59 is Ala or any other amino acid except Pro;

<220>

<221> MISC_FEATURE

<222> (60)..(60)

<223> Xaa at position 60 is Pro, Thr, Asn, or Ser;

<220>

<221> MISC_FEATURE

<222> (61)..(61)

<223> Xaa at position 61 is Leu, or any other amino acid except Pro;

<220>

<221> MISC_FEATURE

<222> (62)..(62)

<223> Xaa at position 62 is Ser or Thr;

<220>

<221> MISC_FEATURE

<222> (63)..(63)

<223> Xaa at position 63 Ser or Asn;

<220>

<21> MISC_FEATURE

<222> (64)..(64)

<223> Xaa at position 64 is Cys or any other amino acid except Pro;

<220>

<221> MISC_FEATURE

<222> (65)..(65)

<223> Xaa at position 65 is Pro, Ser, or Thr;

<220>

<221> MISC_FEATURE

<222> (66)..(66)

<223> Xaa at position 66 is Ser or Thr;

<220>

<221> MISC_FEATURE

<222> (67)..(67)

<223> Xaa at position 67 is Gln or Asn;

<220>

<221> MISC_FEATURE

<222> (68)..(68)

<223> Xaa at position 68 is Ala or any other amino acid except Pro;

<220>

<221> MISC_FEATURE

<222> (69)..(69)

<223> Xaa at position 69 is Leu, Thr, or Ser;

<220>

<221> MISC_FEATURE

<222> (93)..(93)

<223> Xaa at position 93 is Glu or Asn;

<220>

<221> MISC_FEATURE

<222> (94)..(94)

<223> Xaa at position 94 is Gly or any other amino acid except Pro;

<220>

<221> MISC_FEATURE

<222> (95)..(95)

<223> Xaa at position 95 is Ile, Asn, Ser, or Thr;

<220>

<221> MISC_FEATURE

<222> (97)..(97)

<223> Xaa at position 97 is Pro, Ser, Thr, or Asn;

<220>

<221> MISC_FEATURE

<22> (133)..(133)

<223> Xaa at position 133 is Thr or Asn;

<220>

<221> MISC_FEATURE

<222> (134)..(134)

<223> Xaa at position 134 is Gln or any other amino acid except Pro;

<220>

<221> MISC_FEATURE

<222> (135)..(135)

<223> Xaa at position 135 is Gly, Ser, or Thr;

<220>

<221> MISC_FEATURE

<222> (141)..(141)

<223> Xaa at position 141 is Ala or Asn;

<220>

<221> MISC_FEATURE

<222> (142)..(142)

<223> Xaa at position 142 is Ser or any other amino acid except Pro;

<220>

<221> MISC_FEATURE

<222> (143)..(143)

<223> Xaa at position 143 is Ala, Ser, or Thr.

<400> 1

Thr Pro Leu Gly Pro Ala Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys
1 5 10 15

Xaa Leu Glu Gln Val Arg Lys Ile Gln Gly Asp Gly Ala Ala Leu Gln
20 25 30

Glu Lys Leu Cys Xaa Xaa Xaa Lys Leu Cys His Pro Glu Glu Leu Val
35 40 45

Leu Leu Gly His Ser Leu Gly Ile Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
50 55 60

Xaa Xaa Xaa Xaa Xaa Gln Leu Ala Gly Cys Leu Ser Gln Leu His Ser
65 70 75 80

Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Xaa Xaa Xaa Ser
85 90 95

Xaa Glu Leu Gly Pro Thr Leu Asp Thr Leu Gln Leu Asp Val Ala Asp
100 105 110

Phe Ala Thr Thr Ile Trp Gln Gln Met Glu Glu Leu Gly Met Ala Pro
115 120 125

Ala Leu Gln Pro Xaa Xaa Xaa Ala Met Pro Ala Phe Xaa Xaa Xaa Phe
130 135 140

Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser His Leu Gln Ser Phe
145 150 155 160

Leu Glu Val Ser Tyr Arg Val Leu Arg His Leu Ala Gln Pro
165 170

<210> 2

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 2

acccccctgg gccctgccag ctccctgccc cagagcttcc tgctcaagtg gggggacccg
60

ggacggtcga gggacggggg ctccaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgct ccagcggaat ctcgttcact ccttctaggt cccgctaccg
180

cgtcgcgagg tcgagaagct gtgtgccacc tacaagctgt gccaccccca ggagctggtg
240

ctcttcgaca cacggtggat gttcgacacg gtggggctcc tcgaccacct gctcggacac
300

tctctgggca tcccctgggc tcccctgagc agctgcgacg agcctgtgac agaccgtag
360

gggacccgag gggactcgtc gacgcccage caggccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcggt ccgggacgtc gaccgtccga cgaactcggt tgaggtatcg
480

ggccttttcc tctaccaggg gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atggtccccg aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttggac
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct

aaacggtggt ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

aaccagaccg ccatgccggc cttcgcctct gctttccggg acgtcggggt ggtctggcgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggt tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgcgt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

210 211 212 213 220 223

<210> 3

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 3

acccccctgg gccctgccag ctccctgccc cagagcttcc tgotcaagtg gggggaccgc
60

ggacggtcga gggacggggt ctccaaggac gagttgcct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgt ccagcggaat ctcgttcact ctttctaggt cccgctaccg
180

cgtcgcgagg tcgagaagct gtgtgccacc tacaagctgt gccaccccgga ggagctggtg
240

ctcttcgaca cacggtggat gttcgacacg gtggggctcc tcgaccacct gctcggacac
300

ctgggca tccctgggc tccctgagc agctgcgacg agcctgtgac agaccgtag
580

gggacccgag gggactcgtc gacgccagc caggeccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcggc ccgggacgtc gaccgtccga cgaactcggc tgaggtatcg
480

ggccttttcc tctaccaggc gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atggtccccg aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttgagc
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct
720

aaacgggtggg ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

accagggtg ccatgccggc cttcaactct accttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agttgagatg gaagcagcgc cgggcaggag gggtcctggc tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgcgt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 4

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

23> synthetic construct

<400> 4

acccccctgg gccctgccag ctccctgccc cagagcttcc tgctcaagtg gggggacccg
60

ggacggtcga gggacggggg ctcgaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgct ccagcggaa ctcgttccact cttcttaggt cccgctaccg
180

cgtcgagagg tcgagaagct gtgtaacacc accaagctgt gccaccccca ggagctgggtg
240

ctcttcgaca cattgtgggtg gttcgacacg gtggggctcc tcgaccacct gtcgggacac
300

tctctgggca tccctggggc tccctgagc agctgcgacg agcctgtgac agaccgtag
360

gggacccgag gggactcgtc gacgcccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcggt ccgggacgtc gaccgtccga cgaactcggt tgaggatatc
480

ggccttttcc tctaccaggg gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atgggtccccg aggacgtccg ggaccttccc tagaggcccg agttgggtcc caccttgagc
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct
720

aaacgggtggg ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

accagggtg ccatgccggc cttcgctctt gctttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggg tgccctccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgcgt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 5

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 5

acccccctgg gccctgccag ctccctgcc cagagcttcc tgctcaagtg gggggacccg
60

ggacggtcga gggacgggggt ctccaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgct ccagcggaat ctcgttcact ccttctaggt cccgctaccg
180

cgtcgagagg tcgagaagct gtgtgccacc tacaagctgt gccaccccca ggagctgggtg
240

ctcttcgaca cacggtggat gttcgacacg gtggggctcc tcgaccacct gctcggacac
300

tctctgggca tcccctgggc taacactagc agctgcgacg agcctgtgac agaccgtag
360

gggacccgat tggactcctc gacgcccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcggt ccgggacgtc gaccgtccga cgaactcggt tgaggtatcg
480

ggccttttcc totaccaggg gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atgggtcccc aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttggac

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct
720

aaacgggtggt ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

accagggtg ccatgccggc cttcgctct gctttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggt tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgcgt ctttaaggcac cttgccccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 6

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 6

acccccctgg gccctgccag ctccctgcc cagagcttcc tgctcaagtg gggggaccg
60

ggacggtcga gggacgggggt ctcgaaggac gagttcgcct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgt ccagcggaat ctcgttcact ctttctaggt cccgctaccg
180

tcgctgagg tcgagaagct gtgtgccacc tacaagctgt gccaccccga ggagctggtg
210

ctcttcgaca cacggtggat gttcgacacg gtggggctcc tcgaccacct gtcgggacac
300

tctctgggca tcccctgggc tcccctgagc aattgctgac agcctgtgac agacccgtag
360

gggacccgag gggactcggt aacgaccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gctggtcggt ccgggacgtc gaccgtccga cgaactcggt tgaggtatcg
480

ggccttttcc tctaccaggg gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atgggtccccg aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttgagc
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct
720

aaacgggtggt ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

acccagggtg ccatgccggc cttcgcctct gctttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggctcctggt tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggagggtg cgtaccgct ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 7

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 7

acccccctgg gccctgccag ctccctgccc cagagcttcc tgctcaagtg gggggacccg
60

ggacggtcga gggacggggg ctcgaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgct ccagcgggaat ctcgttcact ccttctaggt cccgctaccg
180

cgtcgagagg tcgagaagct gtgtgccacc tacaagctgt gccaccccgaggagctggg
240

ctcttcgaca cacgggtggat gtctgacacg gtggggctcc tcgaccacct gctcggacac
300

tctctgggca tcgttaacgc taccctgagc agctgagc agcctgtgac agaccgtag
360

caattgcat gggactcgtc gacgcccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcggg ccgggacgtc gaccgtccga cgaactcggg tgaggatatg
480

ggccttttcc tctaccaggg gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atgggtcccc aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttgagc
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggccct
720

aaacgggtgg ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

accagggtg ccatgccggc ctctgcctct gctttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggt tgccctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgcgt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 8

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 8

acccccctgg gccctgccag ctccctgccc cagagcttcc tgctcaagtg gggggacccg
60

ggacggtcga gggacggggg ctccaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgct ccagcgggaat ctcgttcaact ccttctaggt cccgctaccg
180

cgtcgcgagg tcgagaagct gtgtgccacc tacaagctgt gccaccccga ggagctggtg
240

ctcttcgaca cacgggtggat gttcgacacg gtgggggtcc tcgaccacct gtcgggacac
300

tctctgggca tcccctgggc tcccctgagc agctgcgacg agcctgtgac agaccgtag
360

gggacccgag gggactcgtc gacgcccagc aacgccaccc agctggcagg ctgcttgagc
420

caactccata gcgggtcggt gcgggtgggtc gaccgtccga cgaactcggt tgaggtatcg

80

ggccttttcc tctaccaggg gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atgggtccccg aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttggac
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct
720

aaacgggtggt ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

accagggtg ccatgccggc ctctgcctct gctttccggg acgtcgggtg ggteccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggt tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgcgt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 9

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 9

acccccctgg gccctgccag ctccctgccc cagagcttcc tgctcaagtg gggggaccgg
60

gacggtcga gggacggggg ctcgaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgct ccagcgggaat ctcgttcact ctttctaggt cccgctaccg
180

cgtcgcgagg tcgagaagct gtgtgccacc tacaagctgt gccaccccca ggagctgggtg
240

ctcttcgaca cacgggtggat gttcgacacg gtggggctcc tcgaccacct gctcggacac
300

tctctgggca tcccctgggc tcccctgagc agctgcgacg agcctgtgac agacccgtag
360

gggacccgag gggactcgtc gacgcccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcggg cggggacgtc gaccgtccga cgaactcggg tgaggtatcg
480

ggccttttcc tctaccaggg gctcctgcag gccctgaacg ggacctcccc ggaaaaggag
540

atggtccccg aggacgtccg ggacttgccc tggaggcccc agttgggtcc caccttggac
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct
720

aaacgggtggg ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

accagggtg ccatgccggc cttcgctctt gctttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggg tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgctt cttaaggcac cttgccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 10

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 10

acccccctgg gccctgccag ctccctgccc cagagcttcc tgctcaagtg gggggacccg
60

ggacggtcga gggacggggg ctccaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgct ccagcgggaat ctcgttcact ccttctaggt cccgctaccg
180

cgtcgagagg tcgagaagct gtgtaacacc accaagctgt gccaccccca ggagctgggtg
240

cttttcgaca cattgtgggtg gttcgacacg gtggggctcc tcgaccacct gctcggacac
300

tctctgggca tcccctgggc tcccctgagc agctgcgacg agcctgtgac agaccgtag
360

gggacccgag gggactcgtc gacgcccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcggt ccgggacgtc gaccgtccga cgaactcggt tgaggtatcg
480

ggccttttcc tctaccaggg gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atgggtcccc aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttgagc
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggccct
720

acgggtggt ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

aaccagaccg ccatgccggc cttcgcctct gctttccggg acgtcggggt ggtctggcgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggt tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgctt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

1044

<210> 11

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 11

acccccctgg gccctgccag ctccctgccc cagagcttcc tgotcaagtg gggggacccg
60

ggacggtcga gggacgggggt ctccaaggac gagttcgctt tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgtt ccagcgggaat ctcgcttact ccttctaggt cccgctaccg
180

cgtcgcgagg tcgagaagct gtgtaacacc accaagctgt gccacccoga ggagctgggtg
240

ctcttcgaca cattgtgggtg gttcgacacg gtggggctcc tcgaccacct gctcggacac
300

tctctgggca tcccctgggc tcccctgagc agctgcgacg agcctgtgac agaccgtag

60

gggacccgag gggactcgtc gacgcccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcggc ccgggacgtc gaccgtccga cgaactcggc tgaggtatcg
480

ggccttttcc totaccaggc gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atgggtccccg aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttggac
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct
720

aaacgggtggt ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

accagggtg ccatgccggc cttcaactct accttcggg acgtcgggtg ggtcccacgg
840

tacggccgga agttgagatg gaagcagcgc cgggcaggag gggctctggt tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgcgt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 12

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

23> synthetic construct

<400> 12

acccccctgg gccctgccag ctccctgccc cagagcttcc tgetcaagtg gggggaccgg
60

ggacgggtcga gggacgggggt ctccaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg ggcagcgcct ccagcggaaat ctogttcact ctttctaggt cccgctaccg
180

cgtcgagagg tcgagaagct gtgtaacacc accaagctgt gccaccccca ggagctgggtg
240

ctcttcgaca cattgtgggtg gttcgacacg gtggggctcc tcgaccacct gctcggacac
300

tctctgggca tcgttaacgc taccctgagc agctgcgacg agcctgtgac agaccgtag
360

caattgcgat gggactcgtc gacgcccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcgggt ccgggacgtc gaccgtccga cgaactcgggt tgaggtatcg
480

ggccttttcc tctaccaggg gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atgggtccccg aggacgtccg ggaccttccc tagaggcccg agttgggtcc caccttggac
600

acactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct
720

aaacgggtggg ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

accagggtg ccatgccggc cttcgctctt gctttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctgggt tgctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ggaggtgt cgtaccgct ctttaaggcac cttgccagc cgcacctcca cagcatggcg
20

cagaattccg tggaacgggt cggg
1044

<210> 13

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 13

acccccctgg gccctgccag ctccctgcc cagagcttcc tgctcaagtg gggggacccg
60

ggacgggtcga gggacgggggt ctccaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgt ccagcggaaat ctcgttcact ccttctaggt cccgctaccg
180

cgtcgagagg tcgagaagct gtgtaacacc accaagctgt gccaccccca ggagctgggtg
240

cttttcgaca cattgtgggtg gttcgacacg gtgggggtcc tcgaccacct gtcgggacac
300

tctctgggca tcccctgggc tcccctgagc agctgagcag agcctgtgac agaccgtag
360

gggacccgag gggactcgtc gacgcccagc aacgccaccc agctggcagg ctgcttgagc
420

caactccata gcgggtcgtt gcgggtgggtc gaccgtccga cgaactcgggt tgaggtatcg
480

ggccttttcc tctaccaggg gtcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atggtccccg aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttgagc
600

actgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggccct
720

aaacgggtgt ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

accagggtg ccatgccggc cttcgectct gctttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggt tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgct ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 14

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 14

acccccctgg gccctgccag ctccctgccc cagagcttcc tgctcaagtg gggggaccog
60

ggacggtcga gggacggggt ctccaaggac gagttcgcct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgct ccagcggaat ctcgttcact ctttctaggt cccgctaccg
180

cgtcgcgagg tcgagaagct gtgtaacacc accaagctgt gccaccccca ggagctggtg

0

ctcttcgaca catttggtg gttcgacacg gtggggctcc tcgaccacct gctcggacac
300

tctctgggca tcccctgggc tcccctgagc agctgcgacg agcctgtgac agacccgtag
360

gggacccgag gggactcgtc gacgcccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gcgggtcggc ccgggacgtc gaccgtccga cgaactcggc tgaggtatcg
480

ggccttttcc tctaccaggc gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540

atgggtccccg aggacgtccg ggaccttccc tagaggaacg gtaccggtcc caccttggac
600

acactgcagc tggacgtcgc cgacttgcca tggccagggt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcctct
720

aaacgggtggt ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780

acccagggtg ccatgccggc cttcgctctt gctttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggt tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggagggtg cgtaccgcgt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 15

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 15

acccccctgg gccctgccag ctccctgccc cagagcttcc tgctcaagtg gggggacccg
60ggacgggtcga gggacgggggt ctccaaggac gagttcgct tagagcaagt gaggaagatc
120cagggcgatg gcgcagcgt ccagcggaaat ctcggttcaact ccttctaggt cccgctaccg
180cgtcgcgagg tcgagaagct gtgtaacacc accaagctgt gccaccccca ggagctgggtg
240ctcttcgaca cattgtgggtg gttcgacacg gtggggctcc tcgaccacct gctcggacac
300tctctgggca tcgttaacgc taccctgagc agctgcgacg agcctgtgac agaccgtag
360caattgcgat gggactcgtc gacgcccagc aacgccaccc agctggcagg ctgcttgagc
420caactccata gcgggtcggt gcgggtgggtc gacggtccga cgaactcgggt tgaggtatcg
480ggccttttcc tctaccaggg gctcctgcag gccctggaag ggatctcccc ggaaaaggag
540atgggtccccg aggacgtccg ggaccttccc tagaggcccc agttgggtcc caccttggac
600aactgcagc tggacgtcgc cgacgggctc aaccagggt ggaacctgtg tgacgtcgac
660ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccc
720aaacggtggt ggtagaccgt cgtctacctt cttgaccctt accggggagc cctgcagccc
780accagggtg ccatgccggc cttegcctct gctttccggg acgtcgggtg ggtcccacgg
840

cgggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggg tgcctcccat
90

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccggt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 16

<211> 1044

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 16

acccccctgg gccctgccag ctccctgccc cagagcttcc tgctcaagtg gggggaccgg
60

ggacgggtcga gggacgggggt ctccaaggac gagttcgct tagagcaagt gaggaagatc
120

cagggcgatg gcgcagcgt ccagcggaaat ctggttcaact cttcttaggt cccgctaccg
180

cgtcgcgagg tcgagaagct gtgtaacacc accaagctgt gccacccga ggagctggg
240

ctcttcgaca cattgtggggt gttcgacacg gtggggctcc tcgaccacct gctcggacac
300

tctctgggca tcccctgggc tcccctgagc aattgcgacg agcctgtgac agaccgtag
360

gggacccgag gggactcgtt aacgaccagc caggccctgc agctggcagg ctgcttgagc
420

caactccata gctgggtcggg ccgggacgtc gaccgtccga cgaactcggg tgaggatatc
480

ggccttttcc tctaccaggg gctcctgcag gccctgaacg ggacctcccc ggaaaaggag
540

atgggtccccg aggacgtccg ggacttgccc tggaggcccc agttgggtcc caccttggaac
600

acactgcagc tggacgtcgc cgacgggctc aaccagggtt ggaacctgtg tgacgtcgac
660

ctgcagcggc tgtttgccac caccatctgg cagcagatgg aagaactggg aatggcccct
720

aaacgggtggt ggtagaccgt cgtctacott cttgaccott accggggagc cctgcagccc
780

accagggttg ccatgccggc cttcgcctct gctttccggg acgtcgggtg ggtcccacgg
840

tacggccgga agcggagacg aaagcagcgc cgggcaggag gggtcctggt tgcctcccat
900

ctgcagagct tcgtcgcggc ccgtcctccc caggaccaac ggagggtaga cgtctcgaag
960

ctggaggtgt cgtaccgctt ctttaaggcac cttgcccagc ccgacctcca cagcatggcg
1020

cagaattccg tggaacgggt cggg
1044

<210> 17

<211> 1762

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 17

gatgcgcaca agagtggagt tgctcatcgg tttaaagatt tgggagaaga aaatttcaaa
60

gccttggtgt tgattgcctt tgctcagtat cttcagcagt gtccatttga agatcatgta

aaattagtga atgaagtaac tgaatttgca aaaacatgtg ttgctgatga gtcagctgaa
180

aattgtgaca aatcacttca tacccttttt ggagacaaat tatgcacagt tgcaactctt
240

cgtgaaacct atggtgaaat ggctgactgc tgtgcaaaac aagaacctga gagaaatgaa
300

tgcttcttgc aacacaaaga tgacaacca aacctcccc gattggtgag accagagggt
360

gatgtgatgt gcaactgcttt tcatgacaat gaagagacat ttttgaaaaa atacttatat
420

gaaattgcca gaagacatcc ttacttttat gccccggaac tccttttctt tgctaaaagg
480

tataaagctg cttttacaga atgttgccaa gctgctgata aagctgcctg cctggtgcca
540

aagctcgatg aacttcggga tgaagggaag gcttcgtctg ccaaacagag actcaagtgt
600

gccagtctcc aaaaatttgg agaaagagct ttcaaagcat gggcagtagc tcgcctgagc
660

cagagatttc ccaaagctga gtttgacaga gtttccaagt tagtgacaga tcttaccaa
720

gtccacacgg aatgctgcca tggagatctg cttgaatgtg ctgatgacag ggcggacctt
780

gccaagtata tctgtgaaaa tcaagattcg atctccagta aactgaagga atgctgtgaa
840

aaacctctgt tggaaaaatc ccactgcatt gccgaagtgg aaaatgatga gatgcctgct
900

gacttgctt cattagctgc tgattttgtt gaaagtaagg atgtttgcaa aaactatgct
960

gaggcaaagg atgtcttcct gggcatgttt ttgtatgaat atgcaagaag gcacctgat
1020

tactctgtcg tgctgctgct gagacttgcc aagacatatg aaaccactct agagaagtgc
1080

tgtgccgctg cagatcctca tgaatgctat gccaaagtgt tcgatgaatt taaacctctt

40

gtggaagagc ctcagaattt aatcaaacia aattgtgagc tttttgagca gcttggagag
1200

tacaaattcc agaatgcgct attagttcgt tacaccaaga aagtacccca agtgtcaact
1260

ccaactcttg tagaggtctc aagaaaccta ggaaaagtgg gcagcaaata ttgtaaacat
1320

cctgaagcaa aaagaatgcc ctgtgcagaa gactatctat ccgtggctct gaaccagtta
1380

tgtgtgttgc atgagaaaac gccagtaagt gacagagtca ccaaatactg cacagaatcc
1440

ttggtgaaca ggcgaccatg cttttcagct ctggaagtcg atgaaacata cgttcccaaa
1500

gagtttaatg ctgaaacatt caccttccat gcagatatat gcacactttc tgagaaggag
1560

agacaaatca agaaacaaac tgcacttggt gagctcgtga aacacaagcc caaggcaaca
1620

aaagagcaac tgaaagctgt tatggatgat ttgcagctt ttgtagagaa gtgctgcaag
1680

gctgacgata aggagacctg ctttgccgag gagggtaaaa aacttggtgc tgcaagtcaa
1740

gctgccttag gcttataatg ac
1762

<210> 18

<211> 232

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 18

1 Ala Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
5 10 15

Ala Pro Glu Lys Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
20 25 30

Lys Asp Thr Lys Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
35 40 45

Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
50 55 60

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
65 70 75 80

Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
85 90 95

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
100 105 110

Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
115 120 125

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
130 135 140

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
145 150 155 160

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
165 170 175

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
180 185 190

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
195 200 205

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
210 215 220

Ser Leu Ser Leu Ser Pro Gly Lys
225 230

<210> 19

<211> 229

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 19

Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Glu Phe
1 5 10 15

Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
20 25 30

Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
35 40 45

Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val
50 55 60

Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser
65 70 75 80

Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu
85 90 95

Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser

100

105

110

Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro
 115 120 125

Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln
 130 135 140

Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala
 145 150 155 160

Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 165 170 175

Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu
 180 185 190

Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser
 195 200 205

Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
 210 215 220

Leu Ser Leu Gly Lys
 225

<210> 20

<211> 585

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 20

P-15648.ST25.txt

Pro Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu
5 10 15

Glu Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln
20 25 30

Gln Cys Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu
35 40 45

Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys
50 55 60

Ser Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu
65 70 75 80

Arg Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro
85 90 95

Glu Arg Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu
100 105 110

Pro Arg Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His
115 120 125

Asp Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg
130 135 140

Arg His Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg
145 150 155 160

Tyr Lys Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala
165 170 175

Cys Leu Leu Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser
180 185 190

Ser Ala Lys Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly Glu
195 200 205

Arg Ala Phe Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe Pro
 210 215 220

Lys Ala Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys
 225 230 235 240

Val His Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp
 245 250 255

Arg Ala Asp Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser
 260 265 270

Ser Lys Leu Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser His
 275 280 285

Cys Ile Ala Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro Ser
 290 295 300

Leu Ala Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr Ala
 305 310 315 320

Glu Ala Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala Arg
 325 330 335

Arg His Pro Asp Tyr Ser Val Val Leu Leu Leu Arg Leu Ala Lys Thr
 340 345 350

Tyr Glu Thr Thr Leu Glu Lys Cys Cys Ala Ala Ala Asp Pro His Glu
 355 360 365

Cys Tyr Ala Lys Val Phe Asp Glu Phe Lys Pro Leu Val Glu Glu Pro
 370 375 380

Gln Asn Leu Ile Lys Gln Asn Cys Glu Leu Phe Glu Asn Leu Gly Glu
 385 390 395 400

Tyr Lys Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val Pro
 405 410 415

Gln Val Ser Thr Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly Lys
420 425 430

Val Gly Ser Lys Cys Cys Lys His Pro Glu Ala Lys Arg Met Pro Cys
435 440 445

Ala Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu Cys Val Leu His
450 455 460

Glu Lys Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser
465 470 475 480

Leu Val Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu Thr
485 490 495

Tyr Val Pro Lys Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala Asp
500 505 510

Ile Cys Thr Leu Ser Glu Lys Glu Arg Gln Ile Lys Lys Gln Thr Ala
515 520 525

Leu Val Glu Leu Val Lys His Lys Pro Lys Ala Thr Lys Glu Gln Leu
530 535 540

Lys Ala Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys
545 550 555 560

Ala Asp Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu Val
565 570 575

Ala Ala Ser Gln Ala Ala Leu Gly Leu
580 585

<210> 21

<211> 703

<212> DNA

13> Artificial Sequence

<220>

<223> synthetic construct

<400> 21

gagcccaaatt cttgtgacaa aactcacaca tgcccaccgt gccagcacc tgaactcctg
60

gggggaccgt cagtcttcct cttcccccca aaaccaagg acaccctcat gatctcccgg
120

accctgagg tcacatgcgt ggtggtggac gtgagccacg aagaccctga ggtcaagttc
180

aactggtacg tggacggcgt ggaggtgcat aatgccaaaga caaagccgcg ggaggagcag
240

tacaacagca cgtaccgtgt ggtcagcgtc ctcaccgtcc tgcaccagga ctgggtgaat
300

ggcaaggagt acaagtgcaa ggtctccaac aaagccctcc cagcccccat cgagaaaacc
360

atctccaaag ccaaagggca gccccgagaa ccacaggtgt acaccctgcc cccatcccgg
420

gaggagatga ccaagaacca ggtcagcctg acctgcctgg tcaaaggctt ctatcccagc
480

gacatcgccg tggagtggga gagcaatggg cagccggaga acaactacaa gaccacgcct
540

cccgtgctgg actccgacgg ctccctcttc ctctatagca agctcaccgt ggacaagagc
600

aggtggcagc aggggaacgt cttctcatgc tccgtgatgc atgaggctct gcacaaccac
660

tacacgcaga agagcctctc cctgtctccg ggtaaattgat agt
703

<210> 22

<211> 981

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 22

tccaccaagg gcccatcggg cttccccgcta gcgccttgct ccaggagcac ctccgagagc
60acagccgccc tgggctgcct ggtcaaggac tacttccccg aaccggtgac ggtgtcgtgg
120aactcaggcg ccctgaccag cggcgtgcac accttccccg ctgtcctaca gtcctcagga
180ctctactccc tcagcagcgt ggtgaccgtg ccctccagca gcttggggcac gaagacctac
240acctgcaacg tagatcaciaa gccagcaac accaagggtg acaagagagt tgagtccaaa
300tatgggtcccc catgcccacc ctgcccagca cctgagttcc tgggggggacc atcagtcttc
360ctgttceccc caaaacccaa ggacactctc atgatctccc ggaccctga ggtcacgtgc
420gtgggtgggtg acgtgagcca ggaagacccc gaggtccagt tcaactggta cgtggatggc
480gtggaggtgc ataatgccaa gacaaagccg cgggaggagc agttcaacag cacgtaccgt
540gtggtcagcg tcctcacctg cctgcaccag gactggctga acggcaagga gtacaagtgc
600aaggtctcca acaaaggcct cccgtcctcc atcgagaaaa ccatctccaa agccaaaggg
660cagccccgag agccacaggt gtacaccctg ccccatccc aggaggagat gaccaagaac
720caggtcagcc tgacctgcct ggtcaaaggc ttctacccca gcgacatcgc cgtggagtgg
780

gagagcaatg ggcagccgga gaacaactac aagaccacgc ctcccggtgct ggactccgac

ggctccttct tctctacag caggctaacc gtggacaaga gcaggtggca ggaggggaat
900

gtcttctcat gctccgtgat gcatgaggct ctgcacaacc actacacaca gaagagcctc
960

tccctgtctc tgggtaaata a
981

<210> 23

<211> 406

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 23

Thr Pro Leu Gly Pro Ala Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys
1 5 10 15

Ala Leu Glu Gln Val Arg Lys Ile Gln Gly Asp Gly Ala Ala Leu Gln
20 25 30

Glu Lys Leu Cys Ala Thr Tyr Lys Leu Cys His Pro Glu Glu Leu Val
35 40 45

Leu Leu Gly His Ser Leu Gly Ile Pro Trp Ala Pro Leu Ser Ser Cys
50 55 60

Pro Ser Gln Ala Leu Gln Leu Ala Gly Cys Leu Ser Gln Leu His Ser
65 70 75 80

Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Glu Gly Ile Ser
85 90 95

```

Pro Glu Leu Gly Pro Thr Leu Asp Thr Leu Gln Leu Asp Val Ala Asp
      100                      105                      110

Phe Ala Thr Thr Ile Trp Gln Gln Met Glu Glu Leu Gly Met Ala Pro
      115                      120                      125

Ala Leu Gln Pro Thr Gln Gly Ala Met Pro Ala Phe Ala Ser Ala Phe
      130                      135                      140

Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser His Leu Gln Ser Phe
      145                      150                      155                      160

Leu Glu Val Ser Tyr Arg Val Leu Arg His Leu Ala Gln Pro Glu Pro
      165                      170                      175

Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu
      180                      185                      190

Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp
      195                      200                      205

Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp
      210                      215                      220

Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly
      225                      230                      235                      240

Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn
      245                      250                      255

Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp
      260                      265                      270

Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro
      275                      280                      285

Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu
      290                      295                      300

```

Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn
305 310 315 320

Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile
325 330 335

Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr
340 345 350

Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys
355 360 365

Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys
370 375 380

Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu
385 390 395 400

Ser Leu Ser Pro Gly Lys
405

<210> 24

<211> 403

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 24

Thr Pro Leu Gly Pro Ala Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys
1 5 10 15

Ala Leu Glu Gln Val Arg Lys Ile Gln Gly Asp Gly Ala Ala Leu Gln

20

25

30

Glu Lys Leu Cys Ala Thr Tyr Lys Leu Cys His Pro Glu Glu Leu Val
 35 40 45

Leu Leu Gly His Ser Leu Gly Ile Pro Trp Ala Pro Leu Ser Ser Cys
 50 55 60

Pro Ser Gln Ala Leu Gln Leu Ala Gly Cys Leu Ser Gln Leu His Ser
 65 70 75 80

Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Glu Gly Ile Ser
 85 90 95

Pro Glu Leu Gly Pro Thr Leu Asp Thr Leu Gln Leu Asp Val Ala Asp
 100 105 110

Phe Ala Thr Thr Ile Trp Gln Gln Met Glu Glu Leu Gly Met Ala Pro
 115 120 125

Ala Leu Gln Pro Thr Gln Gly Ala Met Pro Ala Phe Ala Ser Ala Phe
 130 135 140

Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser His Leu Gln Ser Phe
 145 150 155 160

Leu Glu Val Ser Tyr Arg Val Leu Arg His Leu Ala Gln Pro Glu Ser
 165 170 175

Lys Tyr Gly Pro Pro Cys Pro Ser Cys Pro Ala Pro Glu Phe Leu Gly
 180 185 190

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
 195 200 205

Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln
 210 215 220

Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val
 230 235 240

His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr
 245 250 255

Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
 260 265 270

Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile
 275 280 285

Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
 290 295 300

Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser
 305 310 315 320

Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
 325 330 335

Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 340 345 350

Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val
 355 360 365

Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met
 370 375 380

His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 385 390 395 400

Leu Gly Lys

<210> 25

<211> 500

<12> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 25

Thr Pro Leu Gly Pro Ala Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys
1 5 10 15

Ala Leu Glu Gln Val Arg Lys Ile Gln Gly Asp Gly Ala Ala Leu Gln
20 25 30

Glu Lys Leu Cys Ala Thr Tyr Lys Leu Cys His Pro Glu Glu Leu Val
35 40 45

Leu Leu Gly His Ser Leu Gly Ile Pro Trp Ala Pro Leu Ser Ser Cys
50 55 60

Pro Ser Gln Ala Leu Gln Leu Ala Gly Cys Leu Ser Gln Leu His Ser
65 70 75 80

Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Glu Gly Ile Ser
85 90 95

Pro Glu Leu Gly Pro Thr Leu Asp Thr Leu Gln Leu Asp Val Ala Asp
100 105 110

Phe Ala Thr Thr Ile Trp Gln Gln Met Glu Glu Leu Gly Met Ala Pro
115 120 125

Ala Leu Gln Pro Thr Gln Gly Ala Met Pro Ala Phe Ala Ser Ala Phe
130 135 140

Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser His Leu Gln Ser Phe
145 150 155 160

Leu Glu Val Ser Tyr Arg Val Leu Arg His Leu Ala Gln Pro Gly Gly
 165 170 175

Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ala His
 180 185 190

Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu Glu Asn Phe
 195 200 205

Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln Gln Cys Pro
 210 215 220

Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu Phe Ala Lys
 225 230 235 240

Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys Ser Leu His
 245 250 255

Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu Arg Glu Thr
 260 265 270

Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro Glu Arg Asn
 275 280 285

Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu Pro Arg Leu
 290 295 300

Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His Asp Asn Glu
 305 310 315 320

Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg Arg His Pro
 325 330 335

Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys Arg Tyr Lys Ala
 340 345 350

Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala Ala Cys Leu Leu

355

360

365

Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala Ser Ser Ala Lys
 370 375 380

Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly Glu Arg Ala Phe
 385 390 395 400

Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe Pro Lys Ala Glu
 405 410 415

Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr Lys Val His Thr
 420 425 430

Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp Asp Arg Ala Asp
 435 440 445

Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile Ser Ser Lys Leu
 450 455 460

Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser His Cys Ile Ala
 465 470 475 480

Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro Ser Leu Ala Ala
 485 490 495

Asp Phe Val Glu
 500

<210> 26

<211> 69

<212> DNA

<213> Artificial Sequence

<220>

<23> synthetic construct

<400> 26

gtaagcttgc gtcgacgcta gcggcgcgcc gccatggcgc gacctgccac ccagagcccc
60

atgaagctg

69

<210> 27

<211> 61

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 27

ggggcagggg gctggctggg ccagtgagg tggcttcctg cactgtccag agtgcaactgt
60

g

61

<210> 28

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 28

ggacagtgc ggaagccact ccactgggcc cagccagctc cctgccccag agcttcctg
59

<10> 29

<211> 72

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 29

gaacctcgag gatcctcatt agggctgggc aaggtgcctt aagacgcggt acgacacctc
60

caggaagctc tg

72

<210> 30

<211> 69

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 30

gtaagcttgc gtcgacgcta gcggcgcgcc gccatggccg gacctgccac ccagagcccc
60

atgaagctg

69

<210> 31

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 31

gctctaaggc cttgagcagg aagctctggg gcagggagct cgctgggccc agtggag
57

<210> 32

<211> 53

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 32

gggcccagcg agctccctgc cccagagctt cctgctcaag gccttagagc aag
53

<210> 33

<211> 72

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 33

gaacctcgag gatcctcatt agggctgggc aaggtgcctt aagacgcggt acgacacctc
60

caggaagctc tg

72

<210> 34

<211> 69

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 34

gtaagcttgc gtcgacgcta gcggcgcgcc gccatggccg gacctgccac ccagagcccc
60

atgaagctg
69

<210> 35

<211> 61

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 35

gtccgagcag cactagttcc tcggggtggc acagcttggt ggtgttacac agcttctcct
60

g
61

<210> 36

<211> 66

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 36

ggcgagcgc tccaggagaa gctgtgtaac accaccaagc tgtgccaccc cgaggaacta
60

gtgctg
66

<210> 37

<211> 72

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 37

gaacctcgag gatcctcatt agggctgggc aagggtgcctt aagacgcggt acgacacctc
60

caggaagctc tg
72

<210> 38

<211> 69

<212> DNA

<213> Artificial Sequence

<220>

> 23> synthetic construct

<400> 38

gtaagcttgc gtcgacgcta gcggcgcgcc gccatggccg gacctgccac ccagagcccc
60

atgaagctg

69

<210> 39

<211> 61

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 39

gcccggcgct ggaaagcgct ggcgaaggcc ggcattggcg tctggttggg ctgcagggca
60

g

61

<210> 40

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 40

ggcccctgcc ctgcagccca accagaccgc catgccggcc ttcgccagcg ctttccagcg
60

<210> 41

<211> 72

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 41

gaacctcgag gatcctcatt agggctgggc aaggtgcctt aagacgcggt acgacacctc
60

caggaagctc tg
72

<210> 42

<211> 69

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 42

gtaagcttgc gtcgacgcta gcggcgcgcc gccatggccg gacctgccac ccagagcccc
60

atgaagctg
69

<210> 43

<211> 68

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 43

gcccggcgct ggaaggtaga gttgaaggcc ggcattggcac cctgggtggg ctgaagagca
60

ggggccat
68

<210> 44

<211> 74

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 44

gggaatggcc cctgctcttc agcccaccca gggtgccatg ccggccttca actctacctt
60

ccagcgccgg gcag
74

<210> 45

<211> 72

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<200> 45
 gacctcgag gatcctcatt agggctgggc aaggtgcctt aagacgcggt acgacacctc
 60

caggaagctc tg
 72

<210> 46

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 46

gctagcggcg cgccaccatg
 20

<210> 47

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 47

gctcagggtg gcgttaacga tgcccagaga gtg
 33

<210> 48

<211> 30

<212> DNA

13> Artificial Sequence

<220>

<223> synthetic construct

<400> 48

gggcatcggtt aacgctaccc tgagcagctg
30

<210> 49

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 49

gactcgagga tcctcattag ggctggg
27

<210> 50

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 50

gctagcggcg cgccaccatg gccggacctg ccacccag
38

<210> 51

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 51

caagcagccg gccagctggg tggcgttgct ggggcagctg ctcag
45

<210> 52

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 52

gccccagcaa cgccacccag ctggccgggt gcttgag
37

<210> 53

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct
<400> 53
gactcgagga tcctcattag ggctgggcaa ggtgccttaa gacgcgg
47

<210> 54

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

2350123
3456789
10111213
14151617
18192021
22232425
26272829
30313233
34353637
38394041
42434445
46474849
50515253
54555657
58596061
62636465
66676869
70717273
74757677
78798081
82838485
86878889
90919293
94959697
9899100
101102103
104105106
107108109
110111112
113114115
116117118
119120121
122123124
125126127
128129130
131132133
134135136
137138139
140141142
143144145
146147148
149150151
152153154
155156157
158159160
161162163
164165166
167168169
170171172
173174175
176177178
179180181
182183184
185186187
188189190
191192193
194195196
197198199
200201202
203204205
206207208
209210211
212213214
215216217
218219220
221222223
224225226
227228229
230231232
233234235
236237238
239240241
242243244
245246247
248249250
251252253
254255256
257258259
260261262
263264265
266267268
269270271
272273274
275276277
278279280
281282283
284285286
287288289
290291292
293294295
296297298
299300301
302303304
305306307
308309310
311312313
314315316
317318319
320321322
323324325
326327328
329330331
332333334
335336337
338339340
341342343
344345346
347348349
350351352
353354355
356357358
359360361
362363364
365366367
368369370
371372373
374375376
377378379
380381382
383384385
386387388
389390391
392393394
395396397
398399400
401402403
404405406
407408409
410411412
413414415
416417418
419420421
422423424
425426427
428429430
431432433
434435436
437438439
440441442
443444445
446447448
449450451
452453454
455456457
458459460
461462463
464465466
467468469
470471472
473474475
476477478
479480481
482483484
485486487
488489490
491492493
494495496
497498499
500501502
503504505
506507508
509510511
512513514
515516517
518519520
521522523
524525526
527528529
530531532
533534535
536537538
539540541
542543544
545546547
548549550
551552553
554555556
557558559
560561562
563564565
566567568
569570571
572573574
575576577
578579580
581582583
584585586
587588589
590591592
593594595
596597598
599600601
602603604
605606607
608609610
611612613
614615616
617618619
620621622
623624625
626627628
629630631
632633634
635636637
638639640
641642643
644645646
647648649
650651652
653654655
656657658
659660661
662663664
665666667
668669670
671672673
674675676
677678679
680681682
683684685
686687688
689690691
692693694
695696697
698699700
701702703
704705706
707708709
710711712
713714715
716717718
719720721
722723724
725726727
728729730
731732733
734735736
737738739
740741742
743744745
746747748
749750751
752753754
755756757
758759760
761762763
764765766
767768769
770771772
773774775
776777778
779780781
782783784
785786787
788789790
791792793
794795796
797798799
800801802
803804805
806807808
809810811
812813814
815816817
818819820
821822823
824825826
827828829
830831832
833834835
836837838
839840841
842843844
845846847
848849850
851852853
854855856
857858859
860861862
863864865
866867868
869870871
872873874
875876877
878879880
881882883
884885886
887888889
890891892
893894895
896897898
899900901
902903904
905906907
908909910
911912913
914915916
917918919
920921922
923924925
926927928
929930931
932933934
935936937
938939940
941942943
944945946
947948949
950951952
953954955
956957958
959960961
962963964
965966967
968969970
971972973
974975976
977978979
980981982
983984985
986987988
989990991
992993994
995996997
9989991000
100110021003
100410051006
100710081009
101010111012
101310141015
101610171018
101910201021
102210231024
102510261027
102810291030
103110321033
103410351036
103710381039
104010411042
104310441045
104610471048
104910501051
105210531054
105510561057
105810591060
106110621063
106410651066
106710681069
107010711072
107310741075
107610771078
107910801081
108210831084
108510861087
108810891090
109110921093
109410951096
109710981099
110011011102
110311041105
110611071108
110911101111
111211131114
111511161117
111811191120
112111221123
112411251126
112711281129
113011311132
113311341135
113611371138
113911401141
114211431144
114511461147
114811491150
115111521153
115411551156
115711581159
116011611162
116311641165
116611671168
116911701171
117211731174
117511761177
117811791180
118111821183
118411851186
118711881189
119011911192
119311941195
119611971198
119912001201
120212031204
120512061207
120812091210
121112121213
121412151216
121712181219
122012211222
122312241225
122612271228
122912301231
123212331234
123512361237
123812391240
124112421243
124412451246
124712481249
125012511252
125312541255
125612571258
125912601261
126212631264
126512661267
126812691270
127112721273
127412751276
127712781279
128012811282
128312841285
128612871288
128912901291
129212931294
129512961297
129812991300
130113021303
130413051306
130713081309
131013111312
131313141315
131613171318
131913201321
132213231324
132513261327
132813291330
133113321333
133413351336
133713381339
134013411342
134313441345
134613471348
134913501351
135213531354
135513561357
135813591360
136113621363
136413651366
136713681369
137013711372
137313741375
137613771378
137913801381
138213831384
138513861387
138813891390
139113921393
139413951396
139713981399
140014011402
140314041405
140614071408
140914101411
141214131414
141514161417
141814191420
142114221423
142414251426
142714281429
143014311432
143314341435
143614371438
143914401441
144214431444
144514461447
144814491450
145114521453
145414551456
145714581459
146014611462
146314641465
146614671468
146914701471
147214731474
147514761477
147814791480
148114821483
148414851486
148714881489
149014911492
149314941495
149614971498
149915001501
150215031504
150515061507
150815091510
151115121513
151415151516
151715181519
152015211522
152315241525
152615271528
152915301531
153215331534
153515361537
153815391540
154115421543
154415451546
154715481549
155015511552
155315541555
155615571558
155915601561
156215631564
156515661567
156815691570
157115721573
157415751576
157715781579
158015811582
158315841585
158615871588
158915901591
159215931594
159515961597
159815991600
160116021603
160416051606
160716081609
161016111612
161316141615
161616171618
161916201621
162216231624
162516261627
162816291630
163116321633
163416351636
163716381639
164016411642
164316441645
164616471648
164916501651
165216531654
165516561657
165816591660
166116621663
166416651666
166716681669
167016711672
167316741675
167616771678
167916801681
168216831684
168516861687
168816891690
169116921693
169416951696
169716981699
170017011702
170317041705
170617071708
170917101711
171217131714
171517161717
171817191720
172117221723
172417251726
172717281729
173017311732
173317341735
173617371738
173917401741
174217431744
174517461747
174817491750
175117521753
175417551756
175717581759
176017611762
176317641765
176617671768
176917701771
177217731774
177517761777
177817791780
178117821783
178417851786
178717881789
179017911792
179317941795
179617971798
179918001801
180218031804
180518061807
180818091810
181118121813
181418151816
181718181819
182018211822
182318241825
182618271828
182918301831
183218331834
183518361837
183818391840
184118421843
184418451846
184718481849
185018511852
185318541855
185618571858
185918601861
186218631864
186518661867
186818691870
187118721873
187418751876
187718781879
188018811882
188318841885
188618871888
188918901891
189218931894
189518961897
189818991900
190119021903
190419051906
190719081909
191019111912
191319141915
191619171918
191919201921
192219231924
192519261927
192819291930
193119321933
193419351936
193719381939
194019411942
194319441945
194619471948
194919501951
195219531954
195519561957
195819591960
196119621963
196419651966
196719681969
197019711972
197319741975
197619771978
197919801981
198219831984
198519861987
198819891990
199119921993
199419951996
199719981999
200020012002
200320042005
200620072008
200920102011
201220132014
201520162017
201820192020
202120222023
202420252026
202720282029
203020312032
203320342035
203620372038
203920402041
204220432044
204520462047
204820492050
205120522053
205420552056
205720582059
206020612062
206320642065
206620672068
206920702071
207220732074
207520762077
207820792080
208120822083
208420852086
208720882089
209020912092
209320942095
209620972098
209921002101
210221032104
210521062107
210821092110
211121122113
211421152116
211721182119
212021212122
212321242125
212621272128
212921302131
213221332134
213521362137
213821392140
214121422143
214421452146
214721482149
215021512152
215321542155
215621572158
215921602161
216221632164
216521662167
216821692170
217121722173
217421752176
217721782179
218021812182
218321842185
218621872188
218921902191
219221932194
219521962197
219821992200
220122022203
220422052206
220722082209
221022112212
221322142215
221622172218
221922202221
222222232224
222522262227
222822292230
223122322233
223422352236
223722382239
224022412242
224322442245
224622472248
224922502251
225222532254
225522562257
225822592260
226122622263
226422652266
226722682269
227022712272
227322742275
227622772278
227922802281
228222832284
228522862287
228822892290
229122922293
229422952296
229722982299
230023012302
230323042305
230623072308
230923102311
231223132314
231523162317
231823192320
232123222323
232423252326
232723282329
233023312332
233323342335
233623372338
233923402341
234223432344
234523462347
234823492350
235123522353
235423552356
235723582359
236023612362
236323642365
236623672368
236923702371
237223732374
237523762377
237823792380
238123822383
238423852386
238723882389
239023912392
239323942395
239623972398
239924002401
240224032404
240524062407
240824092410
241124122413
241424152416
241724182419
242024212422
242324242425
242624272428
242924302431
243224332434
243524362437
243824392440
244124422443
244424452446
244724482449
245024512452
245324542455
245624572458
245924602461
246224632464
246524662467
246824

13> Artificial Sequence

<220>

<223> synthetic construct

<400> 56
gctaacctga ctagttgccc cagccag
27

<210> 57

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 57
gactcgagga tcctcattag ggctggg
27

<210> 58

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 58
gctagcggcg cgccaccatg
20

10> 59

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 59

ggtgcaattg ctcaggggag cccag
25

<210> 60

<211> 23

<212> DNA

213 <213> Artificial Sequence

 $\langle 220 \rangle$

<223> synthetic construct

<400> 60

gcaattgcac cagccaggcc ctg
23

<210> 61

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<100> 61
gactcgagga tcctcattag ggctggg
27

<210> 62

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 62
gctagcggcg cgccaccatg gccggacctg ccacccag
38

<210> 63

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 63
ccggactggt cccgttcagg gcctgcagga gcccttg
37

<210> 64

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 64
gaacgggacc agtccggagt tgggtccac cttgg
35

<210> 65

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 65
gactcgagga tcctcattag ggctgggcaa ggtgccttaa gacgcgg
47

<210> 66

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 66
gtcgacgcta gcggcgcgcc accatggccg gacctg
36